

**ANNUAL REPORT**

**Data collection and analysis in support of single and multispecies  
stock assessments in Chesapeake Bay:**

**The Chesapeake Bay Multispecies Monitoring and Assessment Program**

Prepared for:

**Virginia Marine Resources Commission**

and

**U.S. Fish & Wildlife Service**

For Sampling During:

**Calendar Year 2024 and Previous Years**

Project Number:

**F-130-R-20**

Submitted:

**30 May 2025**

Prepared by:

**Gina M. Ralph  
James Gartland  
Debra J. Gauthier  
Jameson Gregg  
Robert J. Latour**

**School of Marine Science  
College of William and Mary  
Virginia Institute of Marine Science  
Gloucester Point, VA 23062**

## Table of Contents

Abstract .....	2
Introduction.....	3
Methods .....	4
Task 1 .....	4
Task 2 .....	6
Task 3 .....	7
Task 4 .....	7
Results .....	8
Task 1 .....	8
Tasks 2-4.....	8
Species profiles.....	10
Atlantic croaker, <i>Micropogonias undulatus</i> .....	10
Black sea bass, <i>Centropristis striata</i> .....	20
Bluefish, <i>Pomatomus saltatrix</i> .....	28
Butterfish, <i>Peprilus triacanthus</i> .....	36
Kingfishes, <i>Menticirrhus</i> spp. ....	41
Northern puffer, <i>Sphoeroides maculatus</i> .....	51
Scup, <i>Stenotomus chrysops</i> .....	58
Spot, <i>Leiostomus xanthurus</i> .....	66
Striped bass, <i>Morone saxatilis</i> .....	75
Summer flounder, <i>Paralichthys dentatus</i> .....	84
Weakfish, <i>Cynoscion regalis</i> .....	92
White perch, <i>Morone americana</i> .....	102
References .....	112
Appendix I - Water quality.....	115
Water temperature .....	115
Salinity.....	122
Dissolved oxygen.....	129
Appendix II - History of ChesMMAP sampling design .....	136
Appendix III - Additional species profiles .....	138
Blue crab, <i>Callinectes sapidus</i> .....	138
Clearnose skate, <i>Rostroraja eglanteria</i> .....	140

## Abstract

The threats affecting living marine resources are diverse, including overfishing, climate change, and pollution. In response to long-term challenges in fisheries management, a more holistic evaluation of the natural and anthropogenic drivers of populations sizes is needed. Ecosystem management (EM), a suite of strategies that incorporate ecosystem considerations into fisheries and ecosystem management, can be difficult to implement in practice. In the Mid-Atlantic, efforts to implement EM are ongoing; one output has been the annual 'State of the Ecosystem Report' for the region, synthesizes available data on a variety of environmental, ecological, and socioeconomic factors. Historically, the data needed for EM in Chesapeake Bay were either incomplete or nonexistent. The Chesapeake Bay Multispecies Monitoring and Assessment Program (ChesMMAP) was developed to assist in filling these data gaps, and ultimately to support Bay-specific species and ecosystem assessment modeling. ChesMMAP is a fishery-independent monitoring survey that uses a bottom trawl designed to sample late juvenile-to-adult fishes in the mainstem of Chesapeake Bay. Since 2002, this program has provided data on relative abundance, length, weight, sex ratio, maturity, age, and trophic interactions for several important fish species that inhabit the Bay seasonally. In this annual progress report, we synthesize available biological data on 12 bony fishes that support local recreational fisheries, including abundance (biomass and number), length- and age-structure, sex ratio, maturity stage, and diet composition. However, in 2019, the survey underwent a major redesign: VIMS took possession of a new research vessel, the *R/V Virginia*, and the survey bottom trawl gear was replaced with a net consistent with the gear used by other regional bottom trawl surveys (i.e., the Northeast Area Monitoring and Assessment Program and Northeast Fisheries Science Center trawl surveys). At this time, the survey stratification was revised and changes were made to the cruise schedule. During 15 calibration cruises (2019-2022), 516 paired-tows were completed, which was deemed sufficient for robust statistical analysis. Species-specific intercalibrations have been conducted by applying log-Gaussian Cox processes to the paired-tow data and modeling the size distribution of the population at each sampling site and the size-structured clustering of fish at small temporal and spatial scales. As of 2024, historical data (2002-2018) have been calibrated such that a complete time series is now available. The ChesMMAP data inform Bay- and coast-wide fisheries management decisions and the broader use of these data in theses, dissertations, and the peer-reviewed literature contributes to a better understanding of the Bay ecosystem.

## Introduction

Living marine resources provide important economic, cultural, and social benefits that are threatened by myriad anthropogenic pressures, including overfishing, climate change, and pollution (Steneck and Pauly, 2019). Despite this diversity of threats, managers typically have direct control over few of the factors affecting populations of exploited species. As a result, fisheries management efforts have focused on single-species approaches, where each species is assessed and managed in isolation. However, over the past several decades, a more holistic evaluation of the natural and anthropogenic drivers of population sizes has been identified as necessary to improve the status and sustainability of fisheries (Link et al., 2020).

A suite of strategies, collectively termed ecosystem management (EM), have been developed to incorporate biotic and abiotic drivers of populations into fisheries management (see Dolan et al., 2016). The terms used to refer to these various strategies exist along a continuum reflecting the scale at which ecosystem-level considerations are included in the management framework (Dolan et al., 2016). Similar to single-species approaches to fisheries management, the ecosystem approach to fisheries management (EAFM) typically produces biological reference points for a single species or stock; the primary difference between the two approaches is the inclusion of environmental, ecological, and/or socioeconomic factors in EAFM (Link, 2002). Ecosystem-based fisheries management (EBFM) extends beyond this focus on an individual species to consider trade-offs across the ecosystem, such that multiple, or even all, fisheries are evaluated together to optimize yields (Link, 2010; Link and Marshak, 2022). At this scale, species-specific biological reference points must be supplemented by additional criteria or indicators. For example, aggregate biological reference points, such as multispecies maximum sustainable yield (Gaichas et al., 2012), and systemic reference points, such as ecosystem overfishing (Link, 2021) can be developed and applied to specific ecosystems and fisheries. Finally, ecosystem-based management (EBM) considers the ecosystem holistically, with three focal concerns: sustainability, ecological health, and inclusion of humans in the ecosystem (Arkema et al., 2006).

Despite calls for further implementation of EM, in practice, environmental drivers and trophic interactions are rarely integrated into short-term fisheries management decisions (i.e., on total allowable catch: Skern-Mauritzen et al., 2016). While many of the perceived obstacles to more holistic approaches to fisheries management have been addressed (Patrick and Link, 2015), region-specific challenges remain in implementation (e.g., Cowan et al., 2012; Link and Marshak, 2019). In the Mid-Atlantic, efforts to implement EM are ongoing (Gaichas et al., 2018), resulting in the development of the first 'State of the Ecosystem' Report for the region in 2017. Updated annually, these reports synthesize available data on a variety of environmental, ecological, and socioeconomic factors (NOAA National Marine Fisheries Service, 2023).

Historically, the data needed for EM in Chesapeake Bay were either incomplete or nonexistent. The Chesapeake Bay Multispecies Monitoring and Assessment Program (ChesMMAP) was developed to assist in filling these data gaps, and ultimately to support Bay-specific species and ecosystem assessment modeling. ChesMMAP is a fishery-independent monitoring survey that uses a bottom trawl designed to sample late juvenile-to-adult fishes in the mainstem of Chesapeake Bay. While no single gear or monitoring program can collect all of the data necessary for all species and assessment approaches, ChesMMAP was designed to maximize the biological and ecological information collected for several recreationally, commercially, and ecologically important species in the Bay. Since 2002, this program has provided data on relative abundance,

length, weight, sex ratio, maturity, age, and trophic interactions for several important fish species that inhabit the Bay seasonally.

The overarching goal of ChesMMAP is to collect and provide the data necessary to support species and ecosystem assessment modeling and ultimately support the development of fisheries ecosystem plans and regional ocean plans for Chesapeake Bay. In 2024, ChesMMAP efforts focused on four key tasks:

- Task 1: Conduct research cruises
- Task 2: Synthesize data for single species analyses
- Task 3: Quantify trophic interactions for multispecies analyses
- Task 4: Estimate abundance

### *Objectives*

1. To conduct ChesMMAP research cruises between June 2024 and March 2025, including two full-Bay cruises, with up to 80 sites distributed throughout the mainstem of Chesapeake Bay in June and September and two half-Bay cruises, with up to 45 sites in the lower Bay (Virginia waters) in November and up to 35 sites in the upper Bay (Maryland waters) in March. This objective is associated with Task 1.
2. To estimate the population-level parameters necessary to conduct single and multispecies stock assessments. Those include (when appropriate), abundance (biomass and number), length- and age-structure, sex ratio, maturity stage, and diet composition. This objective is associated with Tasks 2-4. The focal species include:
  - Atlantic croaker, *Micropogonias undulatus*
  - Black sea bass, *Centropristis striata*
  - Bluefish, *Pomatomus saltatrix*
  - Butterfish, *Peprilus triacanthus*
  - Kingfishes, *Menticirrhus* spp.
  - Northern puffer, *Sphoeroides maculatus*
  - Scup, *Stenotomus chrysops*
  - Spot, *Leiostomus xanthurus*
  - Striped bass, *Morone saxatilis*
  - Summer flounder, *Paralichthys dentatus*
  - Weakfish, *Cynoscion regalis*
  - White perch, *Morone americana*
3. To serve as a sampling platform for other Bay-related studies focused on, for example, fish disease, water quality, habitat mapping, etc. This is an additional objective associated with Task 1.

## **Methods**

### Task 1

#### *Field methods*

In 2024, we conducted four ChesMMAAP cruises:

- A half-Bay cruise (35 sites) in March in the upper Bay (Maryland), focusing on anadromous species (mainly striped bass and white perch) during spawning,
- Two full-Bay cruises (80 sites each) in June and September, when the Bay is fully populated with both resident and summer-migrant species, and
- A second half-Bay cruise (45 sites) in November in the lower Bay (Virginia), focusing on species during their migration out of the Bay.

Sampling locations were selected using a stratified random design prior to each cruise and the order in which sites were sampled depended on weather, tides, and other logistical considerations. Stratification was simplified in 2019 to four latitudinal regions (two in Maryland, two in Virginia) and two depth strata within each region ( $\leq 12.2$  m,  $> 12.2$  m), and this stratification scheme was followed for all cruises conducted in 2024.

We used the *R/V Virginia*, a 28.3 m steel hull vessel with twin diesels tied to a single controllable-pitch propeller and a dynamic positioning system for station-holding, equipped with a “200 x 12 cm” bottom trawl rigged with a 3.8 cm cookie sweep and using Thyboron Type IV 44” trawl doors for all cruises in 2024 (see Appendix II for further details). At each sampling site, a full profile of the water column, including water temperature, salinity, and dissolved oxygen data were recorded electronically at approximately 1 m intervals using the Hach Hydrolab MS5 Sonde and Hydras recording software. The net was deployed with a depth-dependent tow-wire scope ratio of 3:1 to 6:1 and towed along the bottom for 20 minutes with the tidal current at approximately 3.0 knots. If obstructions or other logistical issues forced a tow to be shortened, the tow still provided a representative sample for the site as long as it was at least 10 minutes in duration.

After retrieval of the net, the catch was sorted by species (and modal size-class, where appropriate) and a subsample was taken from each species and size-class for full processing. The data collected from the subsampled specimens included length and weight, as well as sex and maturity stage (determined macroscopically). Stomachs were removed and those containing prey items were preserved on-board for post-cruise examination (see Task 3). Sagittal otoliths and/or scales were collected from select managed species at all sites in the mainstem of the Chesapeake Bay; these structures were removed, labeled and stored for later age determination. Aggregate weights were recorded by species/size-class for all specimens not selected for full processing, and either all or a representative subsample were enumerated and measured for length. Standardized quality control procedures were implemented during and at the conclusion of each research cruise to ensure that the data collected in the field were complete and consistent with properly functioning gear, equipment, and protocols.

### *Diet*

Diets, typically inferred by stomach content analysis, provide essential information on the trophic structure of ecosystems, which can be incorporated into fisheries management frameworks through multispecies models. To date, the inclusion of multispecies models into formal stock assessment processes has been relatively limited; most efforts have supplemented single-species assessment models rather than forming the basis of management advice (see [Karp et al., 2023](#)). In this context, forage fishes are of particular interest, and the transition from a single-species approach to an ecosystem approach for Atlantic menhaden (*Brevoortia tyrannus*) represents a

major step forward in quantitative, ecosystem approaches to tactical fisheries management (Anstead et al., 2021).

Stomach samples collected and preserved in the field were brought back to VIMS and processed following standard diet analysis procedures (Hyslop, 1980). In general, these protocols involved identifying each prey item to the lowest possible taxonomic level and recording counts and wet weights of the various items. Several diet indices can be calculated to identify the main prey types for each species, including percent by weight, percent by number, and percent frequency-of-occurrence.

### *Ageing*

Information on the age distribution of fishes can provide essential information for fisheries management, and accurate ageing is imperative for understanding population dynamics and informing stock assessments (Campana, 2001). Calcified structures, such as whole otoliths, transverse sectioned otoliths, and scales, typically display seasonal growth patterns that are interpreted as annuli. Transverse sectioned otolith methodology has been validated on several different species within the Chesapeake Bay (Barbieri et al., 1994; Lowerre-Barbieri et al., 1994; Sipe and Chittenden, 2001, 2002; Ihde and Chittenden Jr, 2002) and is recommended by the Atlantic States Marine Fisheries Commission (ASMFC). Thus, this approach is preferred for the assignment of accurate ages for ChesMMAP samples.

Sagittal otoliths and scales collected in the field were brought back to VIMS to determine age assignments under a standard set of criteria. As only a subsample of individuals can be aged, ChesMMAP-specific age-length keys (ALKs) were developed (Coggins Jr et al., 2013). This was required due to the multiple annual sampling events (i.e., bi-monthly cruises) and inter-cruise growth. The updated ALKs use year-specific data but in-year cruise data are pooled over two seasons, spring (typically, March - June) and summer (typically, July - November). Once the ALKs were established for each season, all non-aged measured specimens were assigned to length bins, the total number of specimens captured within each length bin at each site was summed (specimens which had been aged remained in the assigned age class), and the season-specific age-at-length proportions was applied to those sums to determine the total number of age-specific fish caught at each site.

### Task 2

For the focal species presented here, the vast majority of ageing structures (i.e., otoliths, scales) have been analyzed. Currently, most of the ageing structures that remain to be processed represent species that are: 1) of relatively minor management interest (e.g., oyster toadfish otoliths); 2) involve significantly different preparation and analysis techniques (e.g., elasmobranch vertebrae); or currently have no accepted processing protocols (e.g., butterfish sampled from inshore waters).

Species-specific data were synthesized to characterize age- and length-frequency distributions across various spatial and temporal scales (e.g., by year, season, or region of the Bay) for each species. When available, sex ratio and maturity data were used to develop sex-specific analyses.

### Task 3

As with the ageing structures, the vast majority of stomach samples have been analyzed. The diet indices were calculated using a cluster sampling estimator, as each tow yields a cluster of fish at each sampling location (Buckel et al., 1999). Although a variety of diet indices can be calculated using ChesMMAP data, we focused on the contribution by weight, as trophic models typically use biomass as the metric of interest (e.g., Pauly et al., 2000; Buchheister and Latour, 2016; Anstead et al., 2021). Specifically, the contribution of each prey type to the diet by weight (% $Q_k$ ) is given by:

$$Q_k = \frac{\sum_{i=1}^n M_i q_{ik}}{\sum_{i=1}^n M_i},$$

where

$$q_{ik} = \frac{w_{ik}}{w_i} * 100,$$

and where  $n$  is the number of clusters (species/size-class combinations) of the predator of interest sampled,  $M_i$  is the number of individuals of this predator species represented in cluster  $i$ ,  $w_i$  is the total weight of all prey items encountered in the stomachs of that predator sampled from cluster  $i$ , and  $w_{ik}$  is the total weight of prey type  $k$  in those stomachs.

For this report, standardized categories of prey types (fishes, crustaceans, molluscs, worms, misc.) have been developed for all ChesMMAP species. Only those specific prey types greater than or equal to 1.0% of the overall diet are shown (unless the entire category is less than 1.0%). All other specific prey are lumped into a category called 'x - other' (x = fishes, molluscs, etc.), which is distinct from unidentified prey types within the category. For the reader's convenience, the color scheme used for all focal species is the same.

These indices can be coupled with the information generated from Task 2 such that age-, length-, and sex-specific diet characterizations can be developed for each species. Characterizing spatial and temporal variability in these diets is also possible using ChesMMAP data.

### Task 4

For each species sampled by ChesMMAP, a variety of relative abundance trends can be generated according to year, season, and location within Chesapeake Bay.

As documented in previous reports, the acquisition of a new, state-of-the-art research vessel (*R/V Virginia*), paired with evidence that the original trawl gear configuration was suboptimal, created an opportunity to rethink and redesign the ChesMMAP survey (Latour et al., 2023). As a result, a full intercalibration of the two survey designs was required to ensure an uninterrupted time-series could be developed. Between June 2019 and November 2022, 15 calibration cruises (516 paired tows) were undertaken outside of regular survey operations to ensure that no side-by-side vessel effects would bias the survey data. These trips were made soon after regularly scheduled ChesMMAP cruises were completed. Side-by-side tows were completed as simultaneously as possible with the two vessels within approximately 0.4 km. The intercalibration of the two vessel/gear combinations was approached by applying log-Gaussian Cox processes to the paired-tow data (following Thygesen et al., 2019). The application of this approach to data from several common species was generally successful (see Latour et al., 2023) and was subsequently applied to all species included in this report.

For this report, we present arithmetic mean abundance indices; this approach was chosen for its consistency with design-based analytical methodology and to complement existing abundance indices throughout the region. Delta-lognormal indices, model-based indices, and other methods of calculating relative abundance are being explored and may replace these indices in future reports, on a species-by-species basis (e.g., [Lo et al., 1992](#)).

## Results

### Task 1

The four cruises conducted in 2024, with 240 sites sampled, required a total of 27 work days to complete. As indicated in previous reports, the change in survey design in 2019 (from five 80-station, full-Bay trips prior to 2019 to two 80-station, full-Bay trips and two half-Bay trips from 2019 to present) resulted in a decrease in the total number of work days each year from approximately 40 with the *R/V Bay Eagle* to 25-28 days with the *R/V Virginia*. However, the slower cruising speed of the *R/V Virginia* (~8 kt) compared to the *R/V Bay Eagle* (11-12 kt) and the much higher catch rates and therefore higher on-board processing times with the new gear configuration resulted in a lower average number of stations completed per day since 2019.

As anticipated, catch rates for most species increased substantially coinciding with use of the new gear configuration. For some species, much of the increase is due to catching a broader size range, especially on the smaller end, but the increase is very large for almost every species.

### Tasks 2-4

For the 12 focal species, we present data summaries in the form of species profiles that address each task element (i.e., single-species stock parameter summarizations, trophic interaction summaries, and estimates of abundance). The profiles are organized first by species, then by type of analysis (task); however, note that the analyses are not labeled with a task number.

For each of the focal species (where sufficient high-quality data are available), the following data are included:

1. A summary table with numbers and biomass captured and measured during each survey year, as well as the numbers of ageing structure and stomach samples preserved and processed.
2. A series of maps showing total biomass at each sampling site, for each cruise during 2024. Note that biomass at each site was standardized by the area swept (i.e., tow distance \* net width) and scaled to 10,000 m<sup>2</sup>, such that the resulting biomass is presented as kg 10,000 m<sup>-2</sup>.
3. Figures of overall area-swept-corrected abundance indices by number and biomass, calculated using arithmetic means.
4. Length-frequency data by year, for sexes combined and separately.
5. Age-frequency distributions by year (for those species where appreciable numbers have been captured and otoliths have been processed) in both histogram and bubble plot format.
6. Diet analyses by weight, using all data collected and analyzed 2002-2024.
7. Estimated per capita consumption by year for species that are frequently encountered in the diet of 4 key predators (striped bass, summer flounder, weakfish, and clearnose skate).

Note that the calibration analyses have been completed (see [Latour et al., 2023](#)) and the data from 2002 until June 2019 (when the survey was conducted aboard the *R/V Bay Eagle*) have been adjusted to reflect *R/V Virginia* units; however, we have retained a color scheme that distinguishes between the two time periods during this transition year. Therefore, throughout these profiles, when relevant, years during which the survey was conducted aboard the *R/V Bay Eagle* and using the old trawl configuration are indicated by blue, whereas years during which the survey was conducted aboard the *R/V Virginia* and using the new trawl configuration are indicated by light yellow.

These analyses represent a subset of the biological and ecological analyses which could be completed using ChesMMAAP data. Stock assessments regularly include data from ChesMMAAP, and these data form the basis of many peer-reviewed manuscripts focusing on Chesapeake Bay fishes.

## Species profiles

### Atlantic croaker, *Micropogonias undulatus*

*Abundance:* Atlantic croaker are frequently collected in high abundance in ChesMMAAP catches (Table 1), with over 305,000 individuals captured since 2002. Atlantic croaker declined from relatively high catches in the mid-2000s to a low in 2018. Since 2019, large numbers of small individuals with relatively low total biomass have dominated the catches, suggesting that the new gear is substantially more efficient at capturing these size classes. In 2024, the total number of Atlantic croaker was the highest in the time series.

As a seasonal resident in Chesapeake Bay, Atlantic croaker are typically rare in spring, most abundant in summer cruises and in the southern regions, and decrease in fall during their migration out of the Bay. A similar pattern occurred in 2024, with the majority of Atlantic croaker taken in September in regions C and D (Figure 1).

Through 2018, relative abundance indices were calculated using data collected during May, July, and September, in regions 4 and 5 from only the mid-depth and deep strata. With the 2019 restratification, data from both depth strata of regions C and D in June and September were used. Now that the intercalibration analyses are complete, trends in abundance (by number and biomass) can be compared more directly. Both relative abundance indices increased in the early part of the time series, with peaks in 2008 and 2013, after which both indices experienced a steady decline (Figure 2). Indices have remained low since 2019, but with a slight increasing trend. Although these trends may be related to large-scale, cyclic patterns in abundance of Atlantic croaker, it may also be linked to a decrease in the utilization of Chesapeake Bay relative to the coastal ocean that has not been documented in the more northern Delaware Bay (Schonfeld et al., 2022).

*Length and age:* In the survey, specimens of Atlantic croaker ranged from about 1.4 to 49.9 cm total length. The size distribution of this species exhibited high interannual variation (Figure 3). Since 2010, the presence of larger (> 30 cm) fish has steadily declined and no individuals > 38 cm total length have been captured. The *R/V Virginia* more efficiently collects small specimens of Atlantic croaker, but the trend towards a more restricted length-frequency distribution has continued through 2024. Males and females exhibited similar patterns in annual length-frequency distributions (Figure 4). The sex ratio in the catches was variable, usually close to 1:1; however, in some years, such as 2004 and 2018, high catches of females resulted in sex ratios of 1.5-3.5:1.

Ages of Atlantic croaker taken by the survey range from 0 to 17 years (Figure 5). However, as reported previously, the ChesMMAAP survey collects some very small specimens that are designated as age -1; these are fish that were spawned in late summer and fall of a given year, prior to the birth date (1 January) set during an Atlantic croaker ageing workshop sponsored by the Atlantic States Marine Fisheries Commission (see Bonzek et al., 2022). While older specimens (> 8 years) were not uncommon prior to about 2010, a truncation of the age structure, similar to that observed in the length distribution, has occurred. Catches in 2024 continue to display this reduction in older fishes, with only 20 individuals greater than age-2 captured since 2020.

*Diet:* In the Bay, Atlantic croakers consumed a variety of benthic polychaetes, which represent nearly 40% of their diet by weight (Figure 6). Other invertebrates, including crustaceans, molluscs, and to a lesser degree echinoderms and sea squirts, accounted for another 20% of the diet, and only a small amount of the diet consisted of other fishes. Nearly 30% of the total diet was

made up of unidentified material (animal or otherwise), highlighting the benthic and opportunistic nature of this species' feeding habits.

*Consumption:* Atlantic croaker are a common prey of a variety of Chesapeake Bay predators and are relatively easy to distinguish in stomach contents due to their morphology and distinctive hard parts. Intact Atlantic croaker found in the stomachs ranged from 4.1 to 285 mm total length ( $n > 400$ ). These intact fish made up slightly over half of the total Atlantic croaker prey identified. Per capita consumption of Atlantic croaker consumption by key predators was highly variable across the time series (Figure 7), with peaks in 2006 and 2012 followed by strong peaks in Atlantic croaker abundance in 2007 and 2013. In recent years, increasing trends in abundance were reflected by increasing trends in consumption.

Table 1: Atlantic croaker sampling rates and preserved specimen analysis status by year. Note shaded rows represent the change in vessel and trawl gear beginning in June 2019.

Year	Number Caught	Biomass Caught (kg)	Presence at Index Stations (%)	Number Measured	Age Specimens	Ages Read	Stomach Specimens	Stomachs Analyzed
2002	12,689	2,834.0	68.6	7,082	1,126	1,126	1,104	95
2003	12,217	2,850.3	83.1	5,721	548	548	542	62
2004	20,394	5,330.5	93.5	8,850	717	717	702	254
2005	13,281	3,184.8	89.6	7,757	716	716	704	261
2006	14,878	3,486.6	86.5	8,904	854	854	834	749
2007	12,678	1,963.6	88.5	5,974	526	526	523	503
2008	6,260	1,031.3	65.8	3,070	480	480	460	454
2009	3,797	523.0	82.7	3,250	369	369	361	358
2010	3,243	454.3	67.9	2,355	322	322	317	309
2011	5,187	605.5	67.9	2,776	322	322	291	287
2012	2,448	152.9	50.0	1,998	312	312	280	269
2013	8,971	655.1	53.8	3,684	282	282	237	229
2014	1,449	143.3	34.6	620	111	111	73	71
2015	1,723	167.4	43.6	1,402	160	160	110	107
2016	919	90.6	33.3	551	113	113	69	69
2017	1,318	92.9	35.9	1,037	247	247	190	187
2018	1,164	51.6	26.9	455	88	88	56	56
2019	11,685	919.7	84.4	5,792	354	354	233	227
2020	34,291	1,816.6	84.4	6,970	303	303	194	190
2021	8,832	552.9	81.1	3,849	316	316	205	203
2022	18,038	989.3	71.1	3,788	270	269	136	131
2023	38,530	1,942.1	80.0	8,118	390	390	247	243
2024	70,633	2,487.8	85.6	9,899	502	488	313	302

Atlantic Croaker  
*Micropogonias undulatus*

Biomass  
(kg/10,000 m<sup>2</sup>)

- ≤ 5.0
- > 5.0 - 5.5
- > 5.5 - 22.9
- > 22.9
- no catch

Depth Strata  
(m)

- ≤ 12.2
- > 12.2




Figure 1: Site-level estimates of biomass (kg 10,000 m<sup>-2</sup>) of Atlantic croaker in 2024.




Figure 2: Indices of abundance for Atlantic croaker, by number and biomass, for all ages combined. The calibrated R/V Bay Eagle data are blue, while the R/V Virginia data are light yellow.




Figure 3: Length-frequency of Atlantic croaker from 2002-2024. The calibrated R/V Bay Eagle data are blue, while the R/V Virginia data are light yellow.




Figure 4: Length-frequency of Atlantic croaker from 2002-2024, by sex (F = female, M = male, U = unknown). The calibrated R/V Bay Eagle data are blue, while the R/V Virginia data are light yellow. The numbers above each plot represent sample sizes.




Figure 5: Atlantic croaker age frequency through time, standardized to 4,800 annual trawl minutes. Note the shaded polygon represents the change in vessel and trawl gear beginning in June 2019.





Figure 6: Diet composition of Atlantic croaker, expressed as percent by weight, based on 5629 fish and 2557 clusters (2002-2024).



*Figure 7: Annual average consumption of Atlantic croaker, expressed as prey biomass (in grams) per predator, based on its occurrence in the stomach contents of 448 individuals of four key predators: striped bass, summer flounder, weakfish, and clearnose skate (2002-2024).*

## Black sea bass, *Centropristis striata*

*Abundance:* Trawling is generally not considered an effective method of sampling structure-oriented species such as black sea bass, and the original ChesMMA gear caught only 2-50 individuals annually (Table 2). Since 2019, with the new gear configuration, catches have been substantially higher (about 200 to over 500 individuals annually). Total catches were relatively variable, with periods of higher and lower catches. Although only 6 years of data are available using the new gear, the variability seems to have continued, with 2024 catching the highest number of black sea bass in the time series.

In 2024, black sea bass were found primarily in the southern regions (C and D), with occasional catches in region B. The highest catches were typically concentrated along the edges of channels (Figure 8).

Through 2018, relative abundance indices were calculated using data collected during July, September, and November, in regions 4 and 5 from all depth strata. With the 2019 restratification, data from both depth strata of regions C and D in June, September, and November were used. Now that the intercalibration analyses are complete, trends in abundance (by number and biomass) can be compared more directly. Both relative abundance indices were quite variable in the early part of the time series (Figure 9); since 2019, indices have been intermediate, though still variable.

*Length and age:* Specimens of black sea bass captured in this survey ranged from about 4 to 27 cm total length. During the early part of the time series, the size distribution exhibited high interannual variation (Figure 10), due in part to the relatively limited number of specimens captured each year. As more specimens have been captured annually since 2019, a more complete picture of the length-frequency distribution of black sea bass in the Bay has been documented. That being said, specimens captured in this survey are generally small relative to the maximum size of this species (61 cm: [Murdy et al., 1997](#)). Due to the small sizes of most individuals captured by ChesMMA, the majority of specimens observed of this protogynous hermaphroditic species have been females (Figure 11). Black sea bass taken by the survey were young, ranging from 0 to 2 years; individuals aged 1 or even 0 dominate the catches in most years (12).

*Diet:* In the Bay, black sea bass primarily consume small-bodied crustaceans (e.g., mysids, amphipods, isopods, mud crabs), which represent over 60% of their diet by weight (Figure 13). Other invertebrates, including polychaetes, brittle stars, and razor clams, account for another 20% of the diet. Only a small amount of the diet consists of other fishes, particularly bay anchovy. Less than 10% of the total diet is made up of unidentified material (animal or otherwise).

*Consumption:* Black sea bass are easily distinguished from other fishes in stomach contents due to their pigmentation and large spines; however, they are rarely encountered in ChesMMA stomach samples of key predators (n = 6). Therefore, patterns in per capita consumption are not provided.

Table 2: Black sea bass sampling rates and preserved specimen analysis status by year. Note shaded rows represent the change in vessel and trawl gear beginning in June 2019.

Year	Number Caught	Biomass Caught (kg)	Presence at Index Stations (%)	Number Measured	Age Specimens	Ages Read	Stomach Specimens	Stomachs Analyzed
2002	50	4.4	9.8	50	48	48	46	46
2003	42	5.0	11.8	42	32	32	31	31
2004	14	2.2	7.8	14	14	14	14	14
2005	13	1.7	5.3	13	13	13	13	12
2006	22	1.7	6.7	22	17	17	16	16
2007	30	1.8	13.6	30	30	30	29	28
2008	34	2.2	5.9	34	28	28	26	25
2009	35	2.0	14.1	35	35	35	35	34
2010	23	0.6	8.9	23	23	23	22	22
2011	23	1.4	9.7	23	23	23	21	21
2012	9	0.4	2.3	9	9	9	8	7
2013	2	0.1	1.5	2	2	2	1	1
2014	11	0.6	3.7	11	11	11	8	8
2015	11	0.5	5.9	11	11	11	9	9
2016	42	2.0	16.3	42	42	42	30	29
2017	35	1.3	7.4	35	34	34	22	22
2018	8	0.4	1.5	8	8	8	4	4
2019	445	11.1	51.1	445	209	209	148	147
2020	507	16.7	60.7	507	256	256	192	189
2021	514	19.7	57.8	514	263	263	179	177
2022	220	7.5	40.7	220	155	155	99	99
2023	577	25.0	62.2	577	295	290	176	167
2024	467	16.2	54.8	456	270	270	142	134




Figure 8: Site-level estimates of biomass (kg 10,000 m<sup>-2</sup>) of black sea bass in 2024.




Figure 9: Indices of abundance for black sea bass, by number and biomass, for all ages combined. The calibrated R/V Bay Eagle data are blue, while the R/V Virginia data are light yellow.




Figure 10: Length-frequency of black sea bass from 2002-2024. The calibrated R/V Bay Eagle data are blue, while the R/V Virginia data are in light yellow.




Figure 11: Length-frequency of black sea bass from 2002-2024, by sex (F = female, M = male, U = unknown). The calibrated R/V Bay Eagle data are blue, while the R/V Virginia data are in light yellow. The numbers above each plot represent sample sizes.




Figure 12: Black sea bass age frequency through time, standardized to 4,800 annual trawl minutes. Note the shaded polygon represents the change in vessel and trawl gear beginning in June 2019.




Figure 13: Diet composition of black sea bass, expressed as percent by weight, based on 1242 fish and 622 clusters (2002-2024).

## Bluefish, *Pomatomus saltatrix*

*Abundance:* Due to the fast-swimming and pelagic nature of bluefish, this species is not considered to be well sampled by ChesMMAAP, though some useful information can be generated from these survey data. The original ChesMMAAP gear caught 8-126 individuals annually (Table 3). Since 2019, with the new gear configuration, catches have been more variable, with time-series high catches occurring in 2020 and 2021 (208 and 247 individuals, respectively); catches in 2024 were third highest in the time series.

Bluefish were caught sporadically throughout the Bay, though abundance is generally highest in southern regions (C and D) and shallow depths (Figure 14). Catches were typically highest late in the year, presumably as the young-of-the-year fish are moving into deeper waters in preparation for migration out of the Bay.

Through 2018, relative abundance indices were calculated using data collected during September and November cruises, in regions 4 and 5 (note that some previous reports used all five regions) from all depth strata. With the 2019 restratification, data from both depth strata of regions C and D in September and November were used. Now that the intercalibration analyses are complete, trends in abundance (by number and biomass) can be compared more directly. Both relative abundance indices were variable prior to 2011, with alternating years of high and low abundance (Figure 15). From 2012 to 2018, indices exhibited an increasing trend. Since 2019, indices remained variable but generally at lower levels. *Length and age:* Most individuals sampled in the survey are less than 35 cm fork length and, due to the small number of specimens captured and to the protracted spawning season of this species, it is difficult to differentiate cohorts in length frequencies (Figure 16). No pattern of sexual differentiation by size has been observed and sex ratios are relatively variable (Figure 17). Nearly all bluefish captured in the survey are age-0 or age-1 individuals, and in most years, the majority are age-0 fish (Figure 18).

*Diet:* Bluefish collected in the survey are highly piscivorous, with almost 90% of the diet constituting small-bodied fishes such as bay anchovy, spot, and Atlantic menhaden (Figure 19). Crustaceans, mainly mysids and sand shrimp, represent most of the remainder of the diet.

*Consumption:* Bluefish are relatively soft-bodied, so they are more difficult to identify in stomach contents than many other species and may be underrepresented. Only 13 instances of bluefish in the diet of the four key predators have been recorded. Therefore, patterns in per capita consumption are not provided.

Table 3: Bluefish sampling rates and preserved specimen analysis status by year. Note shaded rows represent the change in vessel and trawl gear beginning in June 2019.

Year	Number Caught	Biomass Caught (kg)	Presence at Index Stations (%)	Number Measured	Age Specimens	Ages Read	Stomach Specimens	Stomachs Analyzed
2002	34	10.7	1.3	34	34	34	24	23
2003	114	31.7	33.0	114	74	74	63	62
2004	28	10.0	12.9	28	27	27	22	22
2005	108	22.2	22.1	108	71	71	60	60
2006	23	5.5	12.7	23	23	23	17	17
2007	58	18.2	31.8	58	50	50	44	44
2008	52	15.8	6.7	52	27	27	14	13
2009	11	2.3	6.7	11	11	11	9	9
2010	126	20.2	3.3	82	30	30	13	12
2011	8	2.3	5.6	8	8	8	7	6
2012	17	4.0	8.3	17	17	17	12	12
2013	32	5.4	7.9	32	32	32	26	26
2014	44	5.9	16.7	44	39	39	26	25
2015	125	18.5	17.8	125	49	49	28	28
2016	36	9.8	6.7	36	36	36	19	19
2017	40	6.6	7.8	40	31	31	20	20
2018	85	8.4	14.4	85	41	41	24	24
2019	35	6.4	6.7	35	33	33	14	14
2020	208	23.2	27.8	208	97	97	54	53
2021	247	23.9	28.9	247	122	122	81	79
2022	60	12.2	18.9	60	52	52	27	26
2023	170	14.4	22.2	170	84	84	48	47
2024	183	12.0	34.4	183	96	96	42	38

Bluefish  
*Pomatomus saltatrix*

Biomass  
(kg/10,000 m<sup>2</sup>)

- ≤ 0.1
- > 0.1 - 0.4
- > 0.4 - 0.8
- > 0.8
- no catch

Depth Strata  
(m)

- ≤ 12.2
- > 12.2

0 10 20 40 60 km




Figure 14: Site-level estimates of biomass (kg 10,000 m<sup>-2</sup>) of bluefish in 2024.




Figure 15: Indices of abundance for bluefish, by number and biomass, for all ages combined. The calibrated R/V Bay Eagle data are blue, while the R/V Virginia data are light yellow.




Figure 16: Length-frequency of bluefish from 2002-2024. The calibrated R/V Bay Eagle data are blue, while the R/V Virginia data are light yellow.




Figure 17: Length-frequency of bluefish from 2002-2024 by sex (F = female, M = male, U = unknown). The calibrated R/V Bay Eagle data are blue, while the R/V Virginia data are light yellow. The numbers above each plot represent sample sizes.




Figure 18: Bluefish age frequency through time, standardized to 4,800 annual trawl minutes. Note the shaded polygon represents the change in vessel and trawl gear beginning in June 2019.




Figure 19: Diet composition of bluefish, expressed as percent by weight, based on 679 fish and 388 clusters (2002-2024).

### Butterfish, *Peprilus triacanthus*

*Abundance:* Butterfish are moderately abundant in the survey, with several hundred to over 2,600 specimens typically captured during any survey year (Table 4). Since 2019, with the new gear configuration, the total numbers of individuals caught have been high; in 2024, the total number of butterfish was the highest in the time series at over 5,000.

Butterfish were caught almost exclusively in the southern regions (C and D) and shallow depths (Figure 20). Butterfish abundance follows a generally predictable annual pattern, building from near-zero during March, increasing in abundance through the spring and summer (albeit still low), and reaching a maximum during the September and November cruises.

Through 2018, relative abundance indices were calculated using data collected during the September and November cruises, in regions 4 and 5 from all depth strata (note that some previous reports used only the mid-depth strata). With the 2019 restratification, data from both depth strata of regions C and D in September and November were used. Now that the intercalibration analyses are complete, trends in abundance (by number and biomass) can be compared more directly. Butterfish indices decreased in the early part of the time series, but have generally increased since 2019 to levels near those of the early 2000s (Figure 21).

*Length and age:* Annual length-frequency distributions appear to reveal at least two year-classes of varying strength present in the Chesapeake Bay fish during any given year (Figure 22), however this will require further analysis. This program (and others) has found butterfish collected from estuarine areas extremely difficult to age. We are still investigating methods to obtain accurate age determinations from otolith samples.

*Diet:* Analyses of butterfish stomachs early in the program revealed a high percentage of generally unidentifiable gelatinous zooplankton and other unidentifiable items. It was determined that further analyses of butterfish diets were not an efficient use of resources and the decision was made to discontinue preservation and analysis of butterfish stomachs. Thus, no diet data are presented here.

*Consumption:* Butterfish are relatively soft-bodied, so they are more difficult to identify in stomach contents than many other species and may be underrepresented. Only 23 instances of butterfish in the diet of the four key predators have been recorded. Therefore, patterns in per capita consumption are not provided.

Table 4: Butterfish sampling rates and preserved specimen analysis status by year. Note shaded rows represent the change in vessel and trawl gear beginning in June 2019.

Year	Number Caught	Biomass Caught (kg)	Presence at Index Stations (%)	Number Measured	Age Specimens	Ages Read	Stomach Specimens	Stomachs Analyzed
2002	310	18.3	18.7	310	170	0	168	158
2003	1,000	57.4	63.6	1,000	334	0	334	17
2004	1,133	113.4	55.9	1,071	316	0	316	1
2005	693	48.0	57.0	693	294	0	293	0
2006	634	43.7	62.0	634	3	0	1	0
2007	204	18.8	47.7	204	0	0	0	0
2008	318	22.0	37.8	318	2	0	0	0
2009	415	18.7	55.6	415	0	0	0	0
2010	429	21.8	36.7	429	0	0	0	0
2011	366	22.5	44.9	366	0	0	0	0
2012	991	65.3	35.7	991	0	0	0	0
2013	220	9.6	29.2	220	1	0	0	0
2014	409	20.2	36.7	409	0	0	0	0
2015	402	25.6	21.1	402	0	0	0	0
2016	300	23.3	28.9	300	0	0	0	0
2017	408	21.8	36.7	408	0	0	0	0
2018	124	6.7	20.0	124	0	0	0	0
2019	828	39.9	35.6	828	0	0	0	0
2020	2,616	75.6	61.1	1,876	0	0	0	0
2021	1,569	73.9	57.8	1,569	0	0	0	0
2022	1,359	62.0	70.0	1,359	0	0	0	0
2023	5,039	194.5	84.4	3,460	0	0	0	0
2024	3,619	135.9	70.0	1,891	322	292	0	0

Butterfish  
*Peprilus triacanthus*

Biomass  
(kg/10,000 m<sup>2</sup>)

- ≤ 0.3
- > 0.3 - 0.8
- > 0.8 - 2.2
- > 2.2
- no catch

Depth Strata  
(m)

- ≤ 12.2
- > 12.2

0 10 20 40 60 km




Figure 20: Site-level estimates of biomass (kg 10,000 m<sup>-2</sup>) of butterfish in 2024.




Figure 21: Indices of abundance for butterfish, by number and biomass, for all ages combined. The calibrated R/V Bay Eagle data are blue, while the R/V Virginia data are light yellow.




Figure 22: Length-frequency of butterfish from 2002-2024.

### Kingfishes, *Menticirrhus* spp.

The ranges of three closely related species, northern kingfish (*Menticirrhus saxatilis*), southern kingfish (*Menticirrhus americanus*), and Gulf kingfish (*Menticirrhus littoralis*) overlap in Chesapeake Bay. While some specimens are easily separable in the field, many are not. We have therefore adopted the practice of combining all of these specimens into a single category of kingfishes (*Menticirrhus* spp.). This practice is consistent with the manner in which these species are landed and reported in the fishery.

*Abundance:* Kingfishes were moderately abundant in the survey, with the original ChesMMAAP gear catching approximately 100-600 individuals annually (Table 5). Since 2019, with the new gear configuration, catches have been much higher, about 1,500-6,000 per year.

Catches of kingfishes occurred almost exclusively in the southern regions (C and D) and shallow depths (Figure 23). Catches were highest in the warmer months and remained high into November.

Through 2018, relative abundance indices were calculated using data collected during May, July, September, and November cruises, in regions 4 and 5 from all depth strata. With the 2019 restratification, data from both depth strata of regions C and D in June, September, and November were used. Now that the intercalibration analyses are complete, trends in abundance (by number and biomass) can be compared more directly. Both relative abundance indices were increasing through about 2010, followed by a decline from 2011-2015 that brought the indices back to levels observed at the beginning of the time series (Figure 24). Indices declined again from a high in 2016; since 2019, indices have been relatively stable at moderate levels.

*Length and age:* Due to the relatively small number of specimens captured during early survey years and the overlapping sizes-at-age, it is difficult to interpret length-frequency distributions, though at least two cohorts are apparent in many years (Figure 25). No differential growth patterns between male and female kingfishes have been observed (Figure 26).

Specimens between ages 0 and 7 have been captured in the survey, with most individuals being aged 4 or younger (Figure 27). Year-classes of high (e.g., 2002) and low (e.g., 2004) abundance do seem to track through the stock from year to year, which indicates consistent survey sampling and otolith analysis. This species did not fully recruit to the original ChesMMAAP sampling gear until at least age-1 and perhaps even age-2, but the new gear appears to more efficiently capture younger, smaller individuals. As this species is not subjected to regular stock assessments, specimen processing is assigned a lower level of priority and there is currently a backlog of unprocessed otoliths dating to 2012; ages were assigned using the age-length key that was developed based on specimens captured and processed to date.

*Diet:* Kingfishes collected in the survey have a varied diet (Figure 28). About 40% of the diet was composed of small-bodied crustaceans, such as amphipods and shrimp. Other invertebrates, including worms and molluscs, made up another 45% of the diet; the remainder of the diet included lancelets, unidentified material, and a small amount of fishes.

*Consumption:* Kingfishes are relatively easy to distinguish in stomach contents and are relatively common in stomach samples of key predators (n = 55 occurrences). No clear patterns were evident in the annual per capita consumption (Figure 29), but the peak in consumption of kingfishes in 2015 preceded a peak in abundance of kingfishes in 2016.



Table 5: Kingfishes sampling rates and preserved specimen analysis status by year. Note shaded rows represent the change in vessel and trawl gear beginning in June 2019.

Year	Number Caught	Biomass Caught (kg)	Presence at Index Stations (%)	Number Measured	Age Specimens	Ages Read	Stomach Specimens	Stomachs Analyzed
2002	143	18.5	16.4	143	91	91	87	79
2003	68	19.2	13.5	68	55	55	55	50
2004	67	16.0	14.0	67	55	55	50	48
2005	86	15.3	19.3	86	72	72	69	68
2006	120	24.1	26.1	120	94	94	84	83
2007	122	17.7	25.6	122	88	88	78	76
2008	333	62.6	21.7	300	113	113	97	97
2009	195	24.8	36.3	195	152	152	135	134
2010	447	82.5	35.8	447	231	231	206	199
2011	336	55.7	32.4	336	176	175	155	155
2012	148	24.6	25.9	148	114	0	96	92
2013	165	32.1	24.0	165	106	0	77	77
2014	76	14.2	12.8	76	57	0	39	36
2015	156	24.1	19.4	156	112	0	61	60
2016	613	80.1	42.8	613	265	0	166	163
2017	361	55.2	30.6	361	198	0	138	136
2018	239	39.0	37.2	239	167	0	104	104
2019	3,871	435.9	71.9	2,904	331	0	217	213
2020	5,767	579.1	88.1	3,163	282	0	192	191
2021	1,409	188.9	75.6	1,409	264	0	181	179
2022	2,970	351.1	78.5	2,057	358	0	231	229
2023	3,717	427.2	77.0	2,565	338	0	229	228
2024	1,786	223.4	69.6	1,559	256	0	168	163

Kingfish  
*Menticirrhus spp.*

Biomass  
(kg/10,000 m<sup>2</sup>)

- ≤ 1.0
- > 1.0 - 2.9
- > 2.9 - 7.0
- > 7.0
- no catch

Depth Strata  
(m)

- ≤ 12.2
- > 12.2

0 10 20 40 60 km




Figure 23: Site-level estimates of biomass (kg 10,000 m<sup>-2</sup>) of kingfishes in 2024.




Figure 24: Indices of abundance for kingfishes, by number and biomass, for all ages combined. The calibrated R/V Bay Eagle data are blue, while the R/V Virginia data are light yellow.




Figure 25: Length-frequency of kingfishes from 2002-2024. The calibrated R/V Bay Eagle data are blue, while the R/V Virginia data are light yellow.




Figure 26: Length-frequency of kingfishes from 2002-2024 by sex (F = female, M = male, U = unknown). The calibrated R/V Bay Eagle data are blue, while the R/V Virginia data are light yellow. The numbers above each plot represent sample sizes.




Figure 27: Kingfishes age frequency through time, standardized to 4,800 annual trawl minutes; the shaded polygon represents the change in vessel and trawl gear beginning in June 2019. Note that the ages of specimens collected from 2012-present were estimated based on season age-length keys developed from aged specimens collected prior to 2012.




Figure 28: Diet composition of kingfishes expressed as percent by weight, based on 2860 fish and 1324 clusters (2002-2024).




Figure 29: Annual average consumption of kingfishes, expressed as prey biomass (in grams) per predator, based on its occurrence in the stomach contents of 49 individuals of four key predators: striped bass, summer flounder, weakfish, and clearnose skate (2002-2024).

### Northern puffer, *Sphoeroides maculatus*

*Abundance:* Catches of northern puffer varied by an order of magnitude among years, from as few as 41 in 2005 to over 600 in 2011 (Table 6). Catch rates with the new sampling gear were comparable to those in previous years.

Typical patterns of abundance for this species in the survey are minimal numbers in spring and early summer, followed by a peak in abundance during the September and/or November cruises, perhaps as the summer residents are migrating toward offshore wintering grounds. This pattern was also observed in 2024 (Figure 30). Catches were consistently greatest in the southern regions (C and D), though the species can occur in the lower part of region B.

Through 2018, relative abundance indices were calculated using data collected during September and November cruises, in regions 4 and 5 from all depth strata. With the 2019 restratification, data from both depth strata of regions C and D in September and November were used. Now that the intercalibration analyses are complete, trends in abundance (by number and biomass) can be compared more directly; however, as catches in the survey are patchy, estimates of abundance for this species are of unknown reliability. Both relative abundance indices exhibited high intra- and inter-annual variability through 2018 (Figure 31). Indices have been moderate and less variable since 2019.

*Length and age:* Specimens measuring 2.5 to 30.5 cm total length have been captured by the survey, though most individuals measured have been between 10 and 25 cm (Figure 32). The length composition varied year to year, likely as a result of varying year-classes entering and leaving the Bay stock. The new trawl gear may capture some number of smaller specimens than were previously observed in the survey. The largest individuals captured have generally been females but there appears to be no overall pattern of differential growth between sexes (Figure 33).

Northern puffer is not a high-priority species for stock assessments and standard ageing protocols have not been established. Thus, ageing of vertebrae has not yet been attempted, though the preserved vertebrae remain in storage.

*Diet:* The diet of northern puffer is diverse and fairly even; in ChesMMAAP specimens, miscellaneous prey items including unidentified material, molluscs, and crustaceans made up approximately equal parts of the diet (Figure 34). Worms contributed nearly all of the remainder of the diet, with fish tissue contributing only minimally to the diet.

*Consumption:* Northern puffer are easily distinguished from other fishes in stomach contents due to their anatomy; however, they are rarely encountered in ChesMMAAP stomach samples of key predators ( $n = 4$ ). Therefore, patterns in per capita consumption are not provided.

Table 6: Northern puffer sampling rates and preserved specimen analysis status by year. Note shaded rows represent the change in vessel and trawl gear beginning in June 2019.

Year	Number Caught	Biomass Caught (kg)	Presence at Index Stations (%)	Number Measured	Age Specimens	Ages Read	Stomach Specimens	Stomachs Analyzed
2002	231	23.9	20.0	231	177	0	171	156
2003	225	32.5	34.1	225	100	0	92	91
2004	41	6.9	9.7	41	31	0	27	26
2005	131	13.7	25.6	131	84	0	84	83
2006	52	5.5	17.7	52	51	0	48	47
2007	155	19.8	75.0	155	127	0	124	124
2008	90	6.9	21.1	90	78	0	77	77
2009	76	7.2	24.4	76	69	0	68	67
2010	326	54.7	44.4	326	176	0	157	156
2011	614	55.0	50.6	614	247	0	238	236
2012	50	5.3	11.9	50	50	0	41	40
2013	63	4.2	15.7	63	61	0	55	52
2014	49	3.6	12.2	49	39	0	16	16
2015	290	44.1	36.7	290	157	0	54	54
2016	519	65.6	40.0	519	231	0	99	97
2017	231	22.4	25.6	231	148	0	116	116
2018	246	24.5	28.9	246	128	0	87	87
2019	143	13.6	22.2	143	99	0	77	74
2020	80	7.0	35.6	80	54	0	23	23
2021	43	5.0	18.9	43	34	0	23	23
2022	57	3.6	21.1	57	42	0	13	13
2023	127	8.1	35.6	127	63	0	46	46
2024	16	1.3	11.1	16	12	0	7	7




Figure 30: Site-level estimates of biomass (kg 10,000 m<sup>-2</sup>) of northern puffer in 2024.




Figure 31: Indices of abundance for northern puffer, by number and biomass, for all ages combined. The calibrated R/V Bay Eagle data are blue, while the R/V Virginia data are light yellow.



Figure 32: Length-frequency of northern puffer from 2002-2024. The calibrated R/V Bay Eagle data are blue, while the R/V Virginia data are light yellow.




Figure 33: Length-frequency of northern puffer from 2002-2024 by sex (F = female, M = male, U = unknown). The calibrated R/V Bay Eagle data are blue, while the R/V Virginia data are light yellow. The numbers above each plot represent sample sizes.




Figure 34: Diet composition of northern puffer, expressed as percent by weight, based on 1711 fish and 760 clusters (2002-2024).

### Scup, *Stenotomus chrysops*

*Abundance:* Total yearly captures of scup are highly variable, probably as a result of both actual coast-wide abundance and availability to the survey gear (Table 7). Since 2019, with the new gear configuration, the total numbers of scup caught have been generally higher than in the early years; however, in 2024, only 15 individuals were caught.

Survey catches of scup are typically rare during spring through early summer and nearly always reach a peak in September before declining again in November as fish leave the Bay; in 2024, scup were caught only in June and September (Figure 35).

Through 2018, relative abundance indices were calculated using data collected during July, September, and November cruises, in regions 4 and 5 from shallow and mid-depth strata. With the 2019 restratification, data from shallow depth strata of regions C and D in June, September, and November were used. Now that the intercalibration analyses are complete, trends in abundance (by number and biomass) can be compared more directly. Both relative abundance indices were highly variable prior to 2019, but have declined steadily since then (Figure 36).

*Length and age:* Most specimens captured in the survey are less than 20 cm fork length and at least two size classes are apparent in length data (Figure 37). Due to the small size and sexual immaturity of the majority of scup sampled, sex cannot be determined in the field for large numbers of specimens. Sex-specific length frequencies do not display any discernible pattern of differences in sex ratios at size (Figure 38).

Nearly all specimens captured are either age-0 or age-1, so it is difficult to discern whether year-class abundance can be followed through time in age frequency figures (Figure 39). Both the length-frequency and age-frequency distributions were similar before and after the gear change.

*Diet:* The diets of scup collected in the survey were composed primarily of worms and miscellaneous items (including unidentified material), representing over 70% by weight, combined (Figure 40). Crustaceans, including mysids and hermit crabs, and primarily unidentified mollusc meat constitute most of the remainder of the diet, with fish tissue contributing only minimally.

*Consumption:* Scup are relatively easy to identify in stomach contents due to their morphology and large spines; however, they are rarely encountered in ChesMMAP stomach samples of key predators (n = 16). Therefore, patterns in per capita consumption are not provided.

Table 7: Scup sampling rates and preserved specimen analysis status by year. Note shaded rows represent the change in vessel and trawl gear beginning in June 2019.

Year	Number Caught	Biomass Caught (kg)	Presence at Index Stations (%)	Number Measured	Age Specimens	Ages Read	Stomach Specimens	Stomachs Analyzed
2002	107	7.8	10.7	84	40	40	39	34
2003	192	11.1	21.5	192	100	100	99	90
2004	475	26.0	41.5	475	155	155	150	144
2005	674	30.6	21.6	674	86	86	85	82
2006	317	12.7	29.6	317	115	115	112	111
2007	211	6.5	36.7	211	128	128	121	119
2008	56	4.1	12.9	56	42	42	42	42
2009	201	6.6	17.2	201	97	97	92	91
2010	853	29.2	25.0	653	126	126	125	123
2011	72	2.7	23.9	72	56	56	51	51
2012	12	0.4	3.4	12	12	12	12	12
2013	49	1.8	7.6	49	28	28	25	23
2014	63	2.6	6.5	63	26	26	19	19
2015	988	45.6	38.7	988	186	186	88	87
2016	65	2.0	9.7	65	40	40	20	20
2017	25	0.4	4.3	25	20	20	12	12
2018	386	12.2	29.0	386	94	94	58	58
2019	1,126	35.1	40.0	883	196	196	135	135
2020	626	18.7	25.7	626	34	34	23	23
2021	1,135	45.7	28.6	1,135	112	112	59	59
2022	419	19.2	28.6	295	90	90	34	30
2023	15	0.4	6.7	15	14	0	2	2
2024	566	16.1	9.5	132	32	28	8	8

Scup  
*Stenotomus chrysops*

Biomass  
(kg/10,000 m<sup>2</sup>)

- ≤ 0.1
- > 0.1 - 0.3
- > 0.3 - 1.4
- > 1.4
- no catch

Depth Strata  
(m)

- ≤ 12.2
- > 12.2

0 10 20 40 60 km




Figure 35: Site-level estimates of biomass (kg 10,000 m<sup>-2</sup>) of scup in 2024.




Figure 36: Indices of abundance for scup, by number and biomass, for all ages combined. The calibrated R/V Bay Eagle data are blue, while the R/V Virginia data are light yellow.




Figure 37: Length-frequency of scup from 2002-2024.




Figure 38: Length-frequency of scup from 2002-2024, by sex (F = female, M = male, U = unknown). The calibrated R/V Bay Eagle data are blue, while the R/V Virginia data are light yellow. The numbers above each plot represent sample sizes.




Figure 39: Scup age frequency through time, standardized to 4,800 annual trawl minutes; the shaded polygon represents the change in vessel and trawl gear beginning in June 2019. Note that the ages of specimens collected from 2008-2012 were estimated based on season age-length keys developed from aged specimens collected from 2002-2019.




Figure 40: Diet composition of scup, expressed as percent by weight, based on 1375 fish and 575 clusters (2002-2024.)

### Spot, *Leiostomus xanthurus*

*Abundance:* Spot are typically among the most abundant species in the survey. Prior to 2014, 2,000-11,500 spot were captured annually; however, total annual catches declined afterwards to a time-series minimum of only 400 fish in 2015, followed by a moderate increase to around 1,500 individuals annually from 2016-2018. Deployment of the new gear in 2019 resulted in increased annual catches of 16-33 times the average catches using the old gear (Table 8), with the total number of individuals caught in 2024 being the highest in the time series.

This species is typically common throughout all cruises, except for March, and this pattern was observed in 2024 (Figure 41). It occurred throughout the Bay, though catches were highest in the southern regions (C and D).

Relative abundance indices were calculated using data collected from all region and depth strata. Prior to 2019, data from the July, September, and November cruises were used; after the 2019 restratification, data from the June, September, and November cruises were used. Now that the intercalibration analyses are complete, trends in abundance (by number and biomass) can be compared more directly. Both relative abundance indices displayed a general increase in the early years, followed by a steady decline through 2015, but have increased to the present (Figure 42).

*Length and age:* Most specimens captured in the survey were 10-25 cm fork length and at least two size classes were apparent in some years (Figure 43). Sex-specific length frequencies do not display any discernible differences between males and females (Figure 44).

Nearly all fish in the survey are either age-0 or age-1, with the oldest fish (5 total specimens) captured at age-4 (Figure 45). As discussed above, even though the age distribution of this species in Chesapeake Bay is not wide, the relative numbers of smaller vs. larger specimens can vary significantly year to year. This likely represents both changes in relative year-class strength and the numbers and sizes of specimens moving into the Bay each year.

*Diet:* The majority of the diet of this bottom-feeding species comprised unidentified material (animal or otherwise) and unidentified polychaetes (Figure 46).

*Consumption:* Spot are a common prey of a variety of Chesapeake Bay predators. Intact specimens are easily distinguished from Atlantic croaker in stomach contents, but their soft bodies hinder identification in more advanced stages of digestion. Intact spot found in the stomachs ranged from 8.5 to 181.8 mm total length ( $n > 400$ ). These intact fish made up only about one-third of the spot prey identified. Peaks in per capita consumption by key predators (Figure 47) approximately align with or follow peaks in abundance in 2005, 2010, and 2023.

Table 8: Spot sampling rates and preserved specimen analysis status by year. Note shaded rows represent the change in vessel and trawl gear beginning in June 2019.

Year	Number Caught	Biomass Caught (kg)	Presence at Index Stations (%)	Number Measured	Age Specimens	Ages Read	Stomach Specimens	Stomachs Analyzed
2002	3,122	443.2	37.0	3,034	672	672	647	19
2003	4,081	568.8	50.9	3,102	414	395	396	4
2004	4,131	419.6	64.4	4,089	619	619	578	19
2005	11,561	1,011.2	73.3	10,690	1,030	1,030	979	3
2006	7,080	700.4	71.0	6,439	680	656	632	7
2007	5,729	462.8	72.3	5,396	626	626	602	4
2008	6,256	417.5	63.3	5,197	785	785	742	734
2009	5,191	682.6	47.1	3,481	465	449	447	442
2010	6,744	255.3	67.2	6,336	687	687	652	623
2011	2,867	278.0	39.0	2,867	352	352	320	316
2012	2,161	114.5	35.9	1,758	345	345	259	253
2013	4,087	316.0	44.4	3,430	428	428	289	278
2014	939	117.3	23.3	939	188	188	89	88
2015	401	54.0	15.4	401	102	102	11	11
2016	1,059	67.2	27.1	835	167	167	43	40
2017	1,586	116.4	26.8	1,586	213	213	105	102
2018	1,635	77.0	32.7	1,635	204	204	101	98
2019	67,938	3,529.2	78.4	22,694	556	556	229	225
2020	132,547	6,173.8	89.3	34,056	370	370	134	131
2021	73,427	3,428.0	84.9	21,513	686	686	283	275
2022	107,849	5,005.6	87.3	24,987	699	699	236	233
2023	155,169	6,866.3	90.2	23,285	677	677	340	337
2024	138,639	4,005.8	93.2	27,473	829	827	330	317

Spot  
*Leiostomus xanthurus*

Biomass  
(kg/10,000 m<sup>2</sup>)

- ≤ 11.4
- > 11.4 - 26.0
- > 26.0 - 60.7
- > 60.7
- no catch

Depth Strata  
(m)

- ≤ 12.2
- > 12.2

0 10 20 40 60 km




Figure 41: Site-level estimates of biomass (kg 10,000 m<sup>-2</sup>) of spot in 2024.




Figure 42: Indices of abundance for spot, by number and biomass, for all ages combined. The calibrated R/V Bay Eagle data are blue, while the R/V Virginia data are light yellow.




Figure 43: Length-frequency of spot from 2002-2024.




Figure 44: Length-frequency of spot from 2002-2024, by sex (F = female, M = male, U = unknown). The calibrated R/V Bay Eagle data are blue, while the R/V Virginia data are light yellow. The numbers above each plot represent sample sizes.




Figure 45: Spot age frequency through time, standardized to 4,800 annual trawl minutes. Note the shaded polygon represents the change in vessel and trawl gear beginning in June 2019.





Figure 46: Diet composition of spot, expressed as percent by weight, based on 4559 fish and 2203 clusters (2002-2024.)



*Figure 47: Annual average consumption of spot, expressed as prey biomass (in grams) per predator, based on its occurrence in the stomach contents of 472 individuals of four key predators: striped bass, summer flounder, weakfish, and clearnose skate (2002-2024).*

## Striped bass, *Morone saxatilis*

*Abundance:* Striped bass are typically captured in relatively high abundance each year, with almost 1,000 specimens collected on average over the entire time series (Table 9). In 2024, the total number of striped bass caught was the lowest in the time series, with only 56 individuals.

This species exhibits consistency in its intra-annual abundance pattern, with large numbers of spawning migrants captured during the March cruise, followed by lower numbers in summer as the spawners leave the Bay. Fewer captures occur in July and September, and higher numbers are encountered again in November as fish school before leaving the Bay for offshore wintering grounds. This pattern was not as clearly observed in 2024; the vast majority of captures were in northern regions (A and B) during March, with only limited captures in September, and none were caught in June and November (Figure 48).

Now that the intercalibration analyses are complete, trends in abundance (by number and biomass) can be compared more directly. Two sets of abundance indices have been calculated for this species:

1. The spring spawning stock was evaluated using data from the March cruises, including all depth strata of the northern regions (1-3 prior to 2019, and A and B from 2019 to present).
2. The summer residents were evaluated using data from November, including all depth strata in the southern regions (4 and 5 prior to 2019, and C and D, from 2019 to present). Note that this fall index differs from what has previously been presented, as regions A and B are no longer sampled in November; further evaluation of the applicability of this alternative approach may be warranted.

The spring relative abundance index based on numbers caught displayed a general increase in the early years, through about 2005, followed by a steady decline through 2017 (Figure 49). The spring relative abundance index based on biomass followed a similar pattern, though a few years of higher biomass (i.e., 2008 and 2012) interrupted the otherwise steady decline in biomass from about 2005 to 2017. A March cruise was not conducted in 2018 due to a funding shortfall. Since 2019, both indices have displayed a generally decreasing trend.

The two fall indices exhibited a peak in 2005, declined until about 2011, and then experienced a period of relatively higher values until 2019 (Figure 49). The 2019 indices were extremely high, but were followed by several years of low values, at or near time-series lows. However, as mentioned earlier, the fall striped bass index has been affected by the change in the annual sampling schedule, as no samples were or will be collected in regions A and B (Maryland) in November. Further work is needed to evaluate how these changes impact the reliability of the fall index.

*Length and age:* Most specimens captured in the survey were less than 60 cm fork length (Figure 50). Due to the relatively long-lived nature of this species, the varying life history scenarios for different portions of the stock and associated variable growth rates, along with variable young-of-year recruitment, it is difficult to differentiate year-classes within length-frequency distributions. The largest individuals, typically mature females captured during spring spawning, approached 100 cm, while resident male fish were captured up to about 50 cm (Figure 51).

Striped bass captured in the survey were typically less than about age-10, with specimens up to age-20, captured relatively infrequently. Age-frequency diagrams revealed trends in year-class

strength, where high or low abundances recorded during one year tend to follow into succeeding years (Figure 52). These patterns were generally supported by strong and weak year-classes as measured by the Maryland and Virginia young-of-year beach seine surveys (Durell and Weedon, 2022; Buchanan et al., 2023). The most recent years appeared to be exhibiting a contraction of the age distribution, with only 7 older than age-10.

*Diet:* Fishes comprised the largest taxonomic group in the diet of striped bass captured in this survey, with bay anchovy making up over 30% of the diet, and all fishes together representing nearly two-thirds of the diet. Crustaceans, primarily small-bodied taxa such as amphipods, mysids, and mantis shrimp, and a variety of worms, make up another 25% of the diet. Miscellaneous items, primarily unidentified material, and molluscs represent only minimal contributions to the diet.

Results of diet analyses from this study differ appreciably from some previous studies using specimens from Chesapeake Bay (e.g., Walter and Austin, 2003). These differences are likely the result of both sampling methodological differences (the broad temporal and geographic scale of ChesMMA as well as the gear used compared to many studies which were limited in temporal or geographical scale or which used capture methodologies which yield a narrower size range) and analytical differences in calculating percentages in the diet, as results similar to those presented here were obtained by Overton et al. (2009), using a broad geographic scope, large size range of individuals, and similar cluster-sampling methodology.

*Consumption:* Striped bass are relatively easy to identify as prey items; however, they are uncommonly encountered in predator stomachs (n = 8). Therefore, patterns in per capita consumption are not provided.

Table 9: Striped bass sampling rates and preserved specimen analysis status by year. Note shaded rows represent the change in vessel and trawl gear beginning in June 2019.

Year	Number Caught	Biomass Caught (kg)	Presence at Index Stations (%)	Number Measured	Age Specimens	Ages Read	Stomach Specimens	Stomachs Analyzed
2002	495	313.9	11.8	495	337	337	248	230
2003	765	710.1	52.5	765	501	501	367	355
2004	918	668.9	65.5	918	590	590	476	458
2005	2,245	982.4	63.7	1,919	724	724	528	513
2006	911	839.1	56.6	911	535	535	412	407
2007	579	423.4	51.9	579	389	389	246	241
2008	472	476.9	47.5	472	380	380	317	309
2009	315	243.1	31.2	315	198	198	152	149
2010	288	285.4	25.3	288	205	205	147	144
2011	284	224.9	36.7	284	237	237	178	178
2012	935	330.5	35.9	935	257	257	197	196
2013	695	482.3	51.9	695	373	373	259	123
2014	578	355.8	31.2	578	255	255	186	183
2015	718	398.5	26.2	718	319	319	133	132
2016	1,266	530.2	67.5	1,266	534	534	280	278
2017	1,466	829.0	38.0	1,313	426	426	270	267
2018	313	157.2	6.7	313	173	173	100	100
2019	2,559	679.0	31.2	1,134	265	265	200	200
2020	2,201	412.1	50.0	1,432	300	300	137	134
2021	1,881	528.8	33.8	1,400	195	195	104	102
2022	548	277.4	33.8	548	155	155	67	64
2023	56	49.7	13.8	56	46	46	32	32
2024	80	62.5	23.8	80	74	74	44	45

Striped Bass  
*Morone saxatilis*

Biomass  
(kg/10,000 m<sup>2</sup>)

- ≤ 2.1
- > 2.1 - 7.3
- > 7.3 - 22.7
- > 22.7
- no catch

Depth Strata  
(m)

- ≤ 12.2
- > 12.2

0 10 20 40 60 km




Figure 48: Site-level estimates of biomass (kg 10,000 m<sup>-2</sup>) of striped bass in 2024.




Figure 49: Indices of abundance for striped bass, by number and biomass, for all ages combined; Sp = spring (March) only and Fa = fall (November) only. The calibrated R/V Bay Eagle data are blue, while the R/V Virginia data are light yellow.




Figure 50: Length-frequency of striped bass from 2002-2024.




Figure 51: Length-frequency of striped bass from 2002-2024, by sex (F = female, M = male, U = unknown). The calibrated R/V Bay Eagle data are blue, while the R/V Virginia data are light yellow. The numbers above each plot represent sample sizes.




Figure 52: Striped bass age frequency through time, standardized to 4,800 annual trawl minutes. Note the shaded polygon represents the change in vessel and trawl gear beginning in June 2019.




Figure 53: Diet composition of striped bass, expressed as percent by weight, based on 4842 fish and 1839 clusters (2002-2024).

### Summer flounder, *Paralichthys dentatus*

*Abundance:* Summer flounder are a primary target species for the survey, with several hundred individuals caught in most years (Table 10). While total numbers caught with the new sampling gear (2019-2022) were significantly higher than those from the later years with the *R/V Bay Eagle*, they were within the range captured in the earliest survey years (i.e, 2002-2008).

This species is typically increasingly abundant from spring into late fall, with highest catches in September and/or November, and this pattern was observed in 2024 (Figure 54). It was most abundant in the southern regions (C and D), though catches were not uncommon in the northern regions in September.

Relative abundance indices were calculated based on data collected during the September and November cruises. Prior to 2019, all depth strata in regions 4 and 5 were used; after the 2019 restratification, data from both depth strata in regions C and D were used. Now that the intercalibration analyses are complete, trends in abundance (by number and biomass) can be compared more directly. Both relative abundance indices were highly variable in the earlier part of the time-series and exhibited a substantial decline from a peak in 2006 to low, consistent values from 2012 to 2024 (Figure 55). Changes in the abundance of summer flounder in Chesapeake Bay have been linked to a decrease in the utilization of the Bay relative to the coastal ocean that has not been documented in the more northern Delaware Bay ([Schonfeld et al., 2022](#)).

*Length and age:* Summer flounder measuring 20-50 cm total length are most common in the survey catches, but specimens as large as 75 cm have been captured (Figure 56). In several years, a large number of fish under 30 cm were present in the Bay. This species exhibits sexually dimorphic growth patterns ([Dery, 1981](#)); the vast majority of ChesMMAp specimens larger than 35 cm and nearly all individuals larger than 40 cm are females (Figure 57).

Most fish in the survey are age-5 and under, and the oldest fish yet captured are three specimens at age-12. It is more difficult, compared to other species, to follow abundance trends of particular year-classes in successive years after age-2 (Figure 58). This could be the result of differential migration patterns among different sized fish or of fishery preferences and/or regulations. As well as the declining abundance estimates described above, the summer flounder occurring in the Bay appear to have a restricted age distribution in recent years (since about 2007), except in 2022, when individuals up to age-6 were caught.

*Diet:* The diet of summer flounder comprised fishes, particularly bay anchovy and weakfish, and crustaceans, primarily small-bodied mysids and shrimps (Figure 59), together representing more than 90% by weight. No other prey group constituted more than 2% of the diet.

*Consumption:* Summer flounder are distinguishable in stomach contents by their morphology and distinct jaws and teeth. However, few are encountered (n = 13). Therefore, patterns in per capita consumption are not provided.

Table 10: Summer flounder sampling rates and preserved specimen analysis status by year. Note shaded rows represent the change in vessel and trawl gear beginning in June 2019.

Year	Number Caught	Biomass Caught (kg)	Presence at Index Stations (%)	Number Measured	Age Specimens	Ages Read	Stomach Specimens	Stomachs Analyzed
2002	770	430.5	42.7	770	649	649	425	410
2003	563	341.5	63.6	562	441	441	325	316
2004	728	309.7	72.0	728	565	565	377	372
2005	759	386.7	89.5	759	669	669	420	410
2006	932	453.1	88.6	932	755	755	444	430
2007	567	259.1	81.8	563	489	489	317	313
2008	638	280.9	77.8	638	543	543	354	348
2009	393	187.1	66.7	393	369	369	243	239
2010	385	180.0	67.8	385	354	354	215	209
2011	211	125.3	62.9	211	208	208	111	107
2012	92	33.4	31.0	92	91	91	57	52
2013	110	35.7	33.7	110	107	107	51	45
2014	63	16.7	30.0	63	63	63	40	40
2015	129	41.9	35.6	129	127	127	72	72
2016	77	21.8	30.0	77	77	77	40	39
2017	135	35.0	28.9	135	128	128	85	84
2018	105	26.5	15.6	105	96	96	44	44
2019	623	78.7	90.0	623	385	385	220	216
2020	286	42.0	64.4	286	215	215	105	105
2021	267	23.6	67.8	267	185	185	82	80
2022	688	74.2	76.7	688	426	426	201	197
2023	375	30.9	72.2	375	251	251	127	125
2024	305	39.3	50.0	305	248	246	90	85

Summer Flounder  
*Paralichthyes dentatus*

Biomass  
(kg/10,000 m<sup>2</sup>)

- ≤ 0.2
- > 0.2 - 0.6
- > 0.6 - 1.2
- > 1.2
- no catch

Depth Strata  
(m)

- ≤ 12.2
- > 12.2




Figure 54: Site-level estimates of biomass (kg 10,000 m<sup>-2</sup>) of summer flounder in 2024.




Figure 55: Indices of abundance for summer flounder by number and biomass, for all ages combined. The calibrated R/V Bay Eagle data are blue, while the R/V Virginia data are light yellow.




Figure 56: Length-frequency of summer flounder from 2002-2024.




Figure 57: Length-frequency of summer flounder from 2002-2024, by sex (F = female, M = male, U = unknown). The calibrated R/V Bay Eagle data are blue, while the R/V Virginia data are light yellow. The numbers above each plot represent sample sizes.




Figure 58: Summer flounder age frequency through time, standardized to 4,800 annual trawl minutes. Note the shaded polygon represents the change in vessel and trawl gear beginning in June 2019.




Figure 59: Diet composition of summer flounder, expressed as percent by weight, based on 4339 fish and 2087 clusters (2002-2024).

## Weakfish, *Cynoscion regalis*

*Abundance:* Weakfish are among the most abundant species in the survey, with 1,000-3,500 individuals caught through 2010 (Table 11). Overall, total catches were lower from 2011 to 2018, though over 1,000 individuals were captured in a few years. In 2019, with the new gear, catches increased 5- to 10-fold, ranging from about 15,000 to 24,000 annual captures.

This seasonal resident typically migrates into the Bay in late spring, early summer, such that March catches are low, but the remainder of the year is characterized by generally high abundance. Peak catches are usually in September and decline somewhat in November as fish begin their late fall migration out of the Bay. In 2024, weakfish were abundant primarily in southern regions in the June, September, and November cruises, especially at moderate depths (Figure 60).

Prior to 2019, relative abundance indices were calculated using all depth strata in regions 4 and 5 from the July, September, and November cruises; after the 2019 restratification, data from both depth strata in regions C and D were used from the June, September, and November cruises. Now that the intercalibration analyses are complete, trends in abundance (by number and biomass) can be compared more directly. Both relative abundance indices exhibited declines from peaks early in the time series to lows around 2015, followed by an increase (Figure 61). Indices have been low and more stable since 2019. The most recent stock assessment found that the coast-wide stock was depleted ([Commission, 2019](#)), and changes in the abundance of weakfish in Chesapeake Bay have been linked to a decrease in the utilization of the Bay relative to the coastal ocean that has not been documented in the more northern Delaware Bay ([Schonfeld et al., 2022](#)).

*Length and age:* Most weakfish captured by the survey measured 20-35 cm total length, but specimens as large as 61.6 cm have been captured (Figure 62). The length-frequency distribution based on samples collected with the new gear (2019-2024) are similar to those in other survey years. Sex-specific length frequencies do not display any discernible differences between males and females (Figure 63).

With only a few exceptions, most fish captured over 40 cm were sampled during the first two years of the survey (2002 and 2003). Likewise, the age structure of Chesapeake Bay weakfish has compressed over the past several years, with few individuals older than age-2 captured in recent years and almost none older than age-3 (Figure 64). In this survey, each sampling year seems to result in (what appear to be) reasonable numbers of young fish but very few of these specimens are captured in successive years as older fish.

*Diet:* Fishes constituted a majority of prey in the weakfish diet, representing over two-thirds by weight, with bay anchovy the predominant prey item (Figure 65). Crustaceans made up much of the remainder of the diet, with other taxa contributing minimally. Notably, weakfish account for more than 1% of prey in the diet of weakfish. The relatively low percent of Atlantic menhaden observed in the survey stomach samples, when compared to earlier studies, may be due to a combination of the truncation of the size range of weakfish in Chesapeake Bay, the broad geographic and temporal scale of this survey, and the cluster-sampling analytical methodology applied here.

*Consumption:* Weakfish are a soft-bodied species that can often be identified when partially digested by their distinct dentition and otoliths. They are a common prey of a variety of Chesapeake Bay predators. Intact weakfish found in the stomachs ranged from 11.1 to 270 mm

total length ( $n > 200$ ). These intact specimens made up about one-third of the weakfish identified in stomach contents. Notably, a strong peak in weakfish consumption by key predators occurred in 2009 followed by a sharp decline until 2013 (Figure 66). This is consistent with weakfish abundance, which was high from 2009 to 2012 and experienced a sharp decline in 2013 to 2014.

Table 11: Weakfish sampling rates and preserved specimen analysis status by year. Note shaded rows represent the change in vessel and trawl gear beginning in June 2019.

Year	Number Caught	Biomass Caught (kg)	Presence at Index Stations (%)	Number Measured	Age Specimens	Ages Read	Stomach Specimens	Stomachs Analyzed
2002	1,734	304.7	30.3	1,692	803	803	607	583
2003	2,315	400.0	55.9	2,198	707	707	654	640
2004	3,851	561.9	69.5	3,551	1,108	1,108	901	889
2005	2,715	378.5	65.6	2,711	1,119	1,119	918	908
2006	1,476	159.5	60.8	1,462	728	728	561	554
2007	1,214	128.0	55.7	1,210	554	554	439	435
2008	812	83.8	42.2	812	368	368	330	322
2009	873	46.2	60.0	873	478	478	387	384
2010	1,207	76.8	60.7	1,207	607	607	542	530
2011	918	57.5	55.2	918	454	454	323	322
2012	886	72.2	35.7	886	328	328	260	256
2013	301	42.0	28.4	301	187	187	130	128
2014	172	8.6	23.0	172	126	126	72	72
2015	688	51.9	26.7	688	285	285	141	140
2016	1,115	91.2	38.5	1,115	281	281	143	141
2017	943	68.3	36.3	943	335	335	194	191
2018	1,621	61.5	43.7	1,621	273	273	173	172
2019	18,987	1,327.2	80.7	11,355	661	661	387	381
2020	23,685	1,305.0	90.4	10,855	372	372	171	168
2021	16,901	1,044.4	85.9	9,794	467	467	277	273
2022	15,304	740.0	81.5	6,625	466	466	207	206
2023	19,266	843.0	85.9	7,687	460	460	236	229
2024	10,657	686.7	83.7	6,014	403	395	249	237

Weakfish  
*Cynoscion regalis*

Biomass  
(kg/10,000 m<sup>2</sup>)

- ≤ 2.6
- > 2.6 - 6.8
- > 6.8 - 15.3
- > 15.3
- no catch

Depth Strata  
(m)

- ≤ 12.2
- > 12.2

0 10 20 40 60 km




Figure 60: Site-level estimates of biomass (kg 10,000 m<sup>-2</sup>) of weakfish in 2024.




Figure 61: Indices of abundance for weakfish, by number and biomass, for all ages combined. The calibrated R/V Bay Eagle data are blue, while the R/V Virginia data are light yellow.




Figure 62: Length-frequency of weakfish from 2002-2024.




Figure 63: Length-frequency of weakfish from 2002-2024, by sex (F = female, M = male, U = unknown). The calibrated R/V Bay Eagle data are blue, while the R/V Virginia data are light yellow. The numbers above each plot represent sample sizes.




Figure 64: Weakfish age frequency through time, standardized to 4,800 annual trawl minutes. Note the shaded polygon represents the change in vessel and trawl gear beginning in June 2019.





Figure 65: Diet composition of weakfish, expressed as percent by weight, based on 8161 fish and 2835 clusters (2002-2024).



*Figure 66: Annual average consumption of weakfish, expressed as prey biomass (in grams) per predator, based on its occurrence in the stomach contents of 494 individuals of four key predators: striped bass, summer flounder, weakfish, and clearnose skate (2002-2024).*

### White perch, *Morone americana*

*Abundance:* White perch can be extremely abundant in the survey samples in the northern regions of the Bay; over the entire time series, 3,000 to almost 16,000 specimens were collected annually (Table 12).

This species is typically most abundant in shallow strata in the northern regions and is rarely captured at the deepest stations. In 2024, the majority of captures of this species occurred in region A in March (Figure 67); only 269 specimens were caught during the remainder of the year, nearly all of which were taken in region A.

Two sets of abundance indices have been calculated for this species:

1. The spring spawning stock was evaluated using data from the March cruises, including the northern regions (1-3) and the shallow and mid-depth strata (prior to 2019); since 2019, only the shallow depth strata of the northern regions (A and B) were included.
2. The summer residents were evaluated using data from September, including the shallow and mid-depth strata in regions 1 and 2 (prior to 2019) and regions A and B (from 2019 to present). Note that this differs from what has been presented previously, as regions A and B are no longer sampled in November; further evaluation of the applicability of this alternative approach may be warranted.

Now that the intercalibration analyses are complete, trends in abundance (by number and biomass) can be compared more directly. Interestingly, the two sets of abundance indices displayed varying trends in abundance (Figure 68). The spring relative abundance indices increased in the early part of the time series, through about 2006-2007, were generally lower until 2015, then increased through 2018. The fall indices were much more variable year-to-year, and often the fall indices were high when spring indices were low (e.g., in 2009 and 2015). A March cruise was not conducted in 2018 due to a funding shortfall.

Note that these results should be interpreted with caution. The ChesMMAAP survey covers only a portion of the range of the species and catches can be significantly influenced by salinity. Furthermore, due to the cessation of November sampling in the northern regions (A and B), the fall white perch indices should be re-evaluated.

*Length and age:* White perch of sizes greater than approximately 15 cm fork length are well sampled in the survey (Figure 69). Due to the relatively small maximum size, long life, and slow growth rates it is difficult to separate year-classes of this species using length-frequency. Length-frequency distributions based on samples collected with the new gear were similar to those from earlier years, though smaller individuals appeared to be somewhat more efficiently sampled using the near gear. Overall, the survey collected more females than males and females reach a slightly larger maximum size as compared to males (Figure 70).

This species is not well sampled by the survey until approximately age-4; however, past that age, the survey appears to adequately represent all age classes. A small number of age-19 specimens have been captured, but most specimens were younger than age-11. The age distribution appeared to be regulated by the relative success of each year-class. Year-class specific peaks in abundance can be easily followed during successive years in survey samples (e.g., 1993, 1996, 2000, 2003, 2011 year-classes).

*Diet:* Crustaceans constituted about one-third of the diet of white perch captured in this survey, primarily small-bodied taxa such as amphipods, isopods, copepods, and mud crabs (Figure 72). Worms and unidentified material (animal or otherwise) made up about half of the diet. A variety of molluscs (*Macoma* spp. and unidentified tissue) and a small amount of fishes (mostly bay anchovy) contributed the remainder of the diet.

*Consumption:* White perch are relatively easy to identify in stomach contents due to their strong spines and otoliths, and they are moderately common in the diet of key predators (n = 62), especially of striped bass. Per capita consumption by key predators peaked in 2008 and 2016 (Figure 73), while abundance peaked in spring 2007 and fall 2015.

Table 12: White perch sampling rates and preserved specimen analysis status by year. Note shaded rows represent the change in vessel and trawl gear beginning in June 2019.

Year	Number Caught	Biomass Caught (kg)	Presence at Index Stations (%)	Number Measured	Age Specimens	Ages Read	Stomach Specimens	Stomachs Analyzed
2002	6,625	996.6	36.4	4,020	552	552	471	402
2003	3,782	511.5	48.8	1,882	177	168	147	127
2004	11,021	1,727.4	56.9	6,677	356	356	270	267
2005	7,243	843.6	48.8	5,884	429	429	287	280
2006	11,980	1,611.0	47.6	5,899	385	385	263	254
2007	4,915	517.9	71.4	3,194	318	318	277	277
2008	2,924	340.1	46.7	2,360	260	257	227	224
2009	5,130	686.2	35.6	1,749	158	151	126	126
2010	2,996	453.6	34.1	1,627	207	207	158	157
2011	4,619	675.1	27.3	2,392	231	231	177	173
2012	3,737	459.9	31.2	2,423	151	151	111	109
2013	3,249	421.1	51.1	2,469	199	199	109	55
2014	3,208	341.6	46.7	1,844	153	153	94	92
2015	13,708	2,157.4	33.3	4,098	188	188	80	81
2016	7,165	979.5	48.9	2,935	208	208	104	103
2017	7,957	1,113.9	43.2	4,517	159	159	84	80
2018	3,777	522.7	0.0	2,131	102	102	47	46
2019	9,870	888.5	10.2	3,367	129	129	80	80
2020	15,945	1,580.0	40.0	3,128	93	93	43	42
2021	11,614	986.5	42.0	4,298	129	129	69	69
2022	4,909	486.9	52.0	2,114	118	118	63	63
2023	1,889	197.1	34.0	1,044	51	51	41	41
2024	2,873	272.1	30.0	1,366	81	79	42	43

White Perch  
*Morone americana*

- Biomass  
(kg/10,000 m<sup>2</sup>)
- ≤ 9.8
  - > 9.8 - 44.9
  - > 44.9 - 115.0
  - > 115.0
  - no catch

- Depth Strata  
(m)
- ≤ 12.2
  - > 12.2




Figure 67: Site-level estimates of biomass (kg 10,000 m<sup>-2</sup>) of white perch in 2024.





Figure 68: Indices of abundance for white perch, by number and biomass, for all ages combined; Sp = spring (March) only and Fa = fall (September). The calibrated R/V Bay Eagle data are blue, while the R/V Virginia data are light yellow.




Figure 69: Length-frequency of white perch from 2002-2024.




Figure 70: Length-frequency of white perch from 2002-2024, by sex (F = female, M = male, U = unknown). The calibrated R/V Bay Eagle data are blue, while the R/V Virginia data are light yellow. The numbers above each plot represent sample sizes.




Figure 71: White perch age frequency through time, standardized to 4,800 annual trawl minutes. Note the shaded polygon represents the change in vessel and trawl gear beginning in June 2019.




Figure 72: Diet composition of white perch, expressed as percent by weight, based on 3191 fish and 1300 clusters (2002-2024).




Figure 73: Annual average consumption of white perch, expressed as prey biomass (in grams) per predator, based on its occurrence in the stomach contents of 448 individuals of four key predators: striped bass, summer flounder, weakfish, and clearnose skate (2002-2024).

## References

- Anstead, K. A., Drew, K., Chagaris, D., Schueller, A. M., McNamee, J. E., Buchheister, A., et al. (2021). The path to an ecosystem approach for forage fish management: A case study of atlantic menhaden. *Frontiers in Marine Science* 8, 607657.
- Arkema, K. K., Abramson, S. C., and Dewsbury, B. M. (2006). Marine ecosystem-based management: From characterization to implementation. *Frontiers in Ecology and the Environment* 4, 525–532.
- Barbieri, L. R., Chittenden, M. E., and Jones, C. M. (1994). Age, growth, and mortality of Atlantic Croaker, *Micropogonias undulatus*, in the Chesapeake Bay region, with a discussion of apparent geographic changes in population dynamics. *Fishery Bulletin* 92, 1.
- Bonzek, C. F., Gartland, J., Gauthier, D. J., and Latour, R. J. (2022). Annual Report - 2021 Data collection and analysis in support of single and multispecies stock assessments in Chesapeake Bay: The Chesapeake Bay Multispecies Monitoring and Assessment Program. Gloucester Point, VA: Virginia Institute of Marine Science, William & Mary. doi: [10.25773/k7xj-e205](https://doi.org/10.25773/k7xj-e205)
- Buchanan, J. R., Fabrizio, M. C., and Tuckey, T. D. (2023). Estimation of Juvenile Striped Bass Relative Abundance in the Virginia Portion of Chesapeake Bay Annual Progress Report: 2022 - 2023. Virginia Institute of Marine Science, William & Mary. doi: [10.25773/myws-fg17](https://doi.org/10.25773/myws-fg17)
- Buchheister, A., and Latour, R. J. (2016). Dynamic trophic linkages in a large estuarine system—support for supply-driven dietary changes using delta generalized additive mixed models. *Canadian Journal of Fisheries and Aquatic Sciences* 73, 5–17.
- Buckel, J. A., Conover, D. O., Steinberg, N. D., and McKown, K. A. (1999). Impact of age-0 bluefish (*Pomatomus saltatrix*) predation on age-0 fishes in the Hudson River estuary: Evidence for density-dependent loss of juvenile striped bass (*Morone saxatilis*). *Canadian Journal of Fisheries and Aquatic Sciences* 56, 275–287.
- Campana, S. (2001). Accuracy, precision and quality control in age determination, including a review of the use and abuse of age validation methods. *Journal of Fish Biology* 59, 197–242.
- Coggins Jr, L. G., Gwinn, D. C., and Allen, M. S. (2013). Evaluation of age–length key sample sizes required to estimate fish total mortality and growth. *Transactions of the American Fisheries Society* 142, 832–840.
- Commission, A. S. M. F. (2019). Weakfish stock assessment update report. Atlantic States Marine Fisheries Commission Arlington, VA.
- Cowan, J. H., Rice, J. C., Walters, C. J., Hilborn, R., Essington, T. E., Day Jr, J. W., et al. (2012). Challenges for implementing an ecosystem approach to fisheries management. *Marine and Coastal Fisheries* 4, 496–510.
- Dery, L. (1981). Post workshop age and growth study of young summer flounder.
- Dolan, T. E., Patrick, W. S., and Link, J. S. (2016). Delineating the continuum of marine ecosystem-based management: A US fisheries reference point perspective. *ICES Journal of Marine Science* 73, 1042–1050.

Durell, E. Q., and Weedon, C. (2022). Striped Bass Seine Survey Juvenile Index Web Page. Available at: [dnr.maryland.gov/fisheries/pages/juvenile-index.aspx](http://dnr.maryland.gov/fisheries/pages/juvenile-index.aspx) (Accessed May 31, 2023).

Gaichas, S. K., DePiper, G. S., Seagraves, R. J., Muffley, B. W., Sabo, M. G., Colburn, L. L., et al. (2018). Implementing ecosystem approaches to fishery management: Risk assessment in the US Mid-Atlantic. *Frontiers in Marine Science* 5, 442.

Gaichas, S., Gamble, R., Fogarty, M., Benoît, H., Essington, T., Fu, C., et al. (2012). Assembly rules for aggregate-species production models: Simulations in support of management strategy evaluation. *Marine Ecology Progress Series* 459, 275–292.

Hyslop, E. (1980). Stomach contents analysis—a review of methods and their application. *Journal of Fish Biology* 17, 411–429.

Ihde, T. F., and Chittenden Jr, M. E. (2002). Comparison of calcified structures for aging spotted seatrout. *Transactions of the American Fisheries Society* 131, 634–642.

Karp, M. A., Link, J. S., Grezlik, M., Cadrin, S., Fay, G., Lynch, P., et al. (2023). Increasing the uptake of multispecies models in fisheries management. *ICES Journal of Marine Science* 80, 243–257.

Latour, R. J., Gartland, J., and Bonzek, C. F. (2023). Motivation, design, and redesign of a bottom trawl survey in Chesapeake Bay, USA. *Frontiers in Marine Science* 10.

Link, J. (2010). *Ecosystem-based fisheries management: Confronting tradeoffs*. Cambridge University Press.

Link, J. S. (2002). What does ecosystem-based fisheries management mean. *Fisheries* 27, 18–21.

Link, J. S. (2021). Evidence of ecosystem overfishing in US large marine ecosystems. *ICES Journal of Marine Science* 78, 3176–3201.

Link, J. S., Huse, G., Gaichas, S., and Marshak, A. R. (2020). Changing how we approach fisheries: A first attempt at an operational framework for ecosystem approaches to fisheries management. *Fish and Fisheries* 21, 393–434.

Link, J. S., and Marshak, A. R. (2019). Characterizing and comparing marine fisheries ecosystems in the United States: Determinants of success in moving toward ecosystem-based fisheries management. *Reviews in Fish Biology and Fisheries* 29, 23–70.

Link, J. S., and Marshak, A. R. (2022). *Ecosystem-based fisheries management: Progress, importance, and impacts in the united states*. Oxford University Press.

Lo, N. C., Jacobson, L. D., and Squire, J. L. (1992). Indices of relative abundance from fish spotter data based on delta-lognormal models. *Canadian Journal of Fisheries and Aquatic Sciences* 49, 2515–2526.

Lowerre-Barbieri, S. K., Chittenden Jr, M. E., and Jones, C. M. (1994). A comparison of a validated otolith method to age weakfish, *Cynoscion regalis*, with the traditional scale method. *Fishery Bulletin* 92.

Murdy, E. O., Birdsong, R. S., and Musick, J. A. (1997). *Fishes of Chesapeake Bay*. Smithsonian Institution Press.

- NOAA National Marine Fisheries Service (2023). State of the Ecosystem 2023: Mid-Atlantic. Mid-Atlantic Fishery Management Council.
- Overton, A. S., Margraf, F. J., and May, E. B. (2009). Spatial and temporal patterns in the diet of striped bass in Chesapeake Bay. *Transactions of the American Fisheries Society* 138, 915–926.
- Patrick, W. S., and Link, J. S. (2015). Myths that continue to impede progress in ecosystem-based fisheries management. *Fisheries* 40, 155–160.
- Pauly, D., Christensen, V., and Walters, C. (2000). Ecopath, ecosim, and ecospace as tools for evaluating ecosystem impact of fisheries. *ICES Journal of Marine Science* 57, 697–706.
- Schonfeld, A. J., Gartland, J., and Latour, R. J. (2022). Spatial differences in estuarine utilization by seasonally resident species in Mid-Atlantic Bight, USA. *Fisheries Oceanography* 31, 615–628.
- Sipe, A. M., and Chittenden, M. E. (2001). A comparison of calcified structures for aging summer flounder, *Paralichthys dentatus*. *Fishery Bulletin* 99, 628.
- Sipe, A. M., and Chittenden, M. E. (2002). A comparison of calcified structures for aging bluefish in the Chesapeake Bay region. *Transactions of the American Fisheries Society* 131, 783–790.
- Skern-Mauritzen, M., Ottersen, G., Handegard, N. O., Huse, G., Dingsør, G. E., Stenseth, N. C., et al. (2016). Ecosystem processes are rarely included in tactical fisheries management. *Fish and Fisheries* 17, 165–175.
- Steneck, R. S., and Pauly, D. (2019). Fishing through the anthropocene. *Current Biology* 29, R987–R992.
- Thygesen, U. H., Kristensen, K., Jansen, T., and Beyer, J. E. (2019). Intercalibration of survey methods using paired fishing operations and log-Gaussian Cox processes. *ICES Journal of Marine Science* 76, 1189–1199.
- Walter, J. F., and Austin, H. M. (2003). Diet composition of large striped bass (*Morone saxatilis*) in Chesapeake Bay. *Fishery Bulletin* 101, 414.

## Appendix I - Water quality

### Water temperature

#### *Interpolations*

Bottom  
Water  
Temperature  
(°C)





Figure 74: Interpolated bottom water temperature for 2024, by cruise.




Figure 75: Interpolated bottom water temperature averaged over 2002 through 2024, by cruise.






Figure 76: Interpolated 2024 bottom water temperature deviations from average, by cruise.

Profiles

**ChesMMAP March 2024 Temperature Profiles**



Region A



Region B







Figure 77: Interpolated bottom water temperature profile for March 2024.


### ChesMMAP June 2024 Temperature Profiles




Region A



Region B



Region C



Region D





Figure 78: Interpolated bottom water temperature profile for June 2024.

### ChesMMAP September 2024 Temperature Profiles





Figure 79: Interpolated bottom water temperature profile for September 2024.

### ChesMMAP November 2024 Temperature Profiles





Figure 80: Interpolated bottom water temperature profile for November 2024.

# Salinity

## Interpolations





Figure 81: Interpolated bottom salinity for 2024, by cruise.



*Figure 82: Interpolated bottom salinity averaged over 2002 through 2024, by cruise.*






Figure 83: Interpolated 2024 bottom salinity deviations from average, by cruise.

Profiles

**ChesMMAP March 2024 Salinity Profiles**



Region A



Region B





Figure 84: Interpolated bottom salinity profile for March 2024.

### ChesMMAP June 2024 Salinity Profiles







Figure 85: Interpolated bottom salinity profile for June 2024.


### ChesMMAP September 2024 Salinity Profiles




Region A



Region B



Region C



Region D





Figure 86: Interpolated bottom salinity profile for September 2024.

### ChesMMAP November 2024 Salinity Profiles





Figure 87: Interpolated bottom salinity profile for November 2024.

## Dissolved oxygen

### Interpolations




Figure 88: Interpolated bottom dissolved oxygen for 2024, by cruise.




Figure 89: Interpolated bottom dissolved oxygen averaged over 2002 through 2024, by cruise.






Figure 90: Interpolated 2024 bottom dissolved oxygen deviations from average, by cruise.

Profiles

ChesMMAP March 2024 Dissolved Oxygen Profiles



Region A



Region B






Figure 91: Interpolated bottom dissolved oxygen profile for March 2024.


### ChesMMAP June 2024 Dissolved Oxygen Profiles




Region A



Region B



Region C



Region D






Figure 92: Interpolated bottom dissolved oxygen profile for June 2024.


### ChesMMAP September 2024 Dissolved Oxygen Profiles




Region A



Region B



Region C



Region D





Figure 93: Interpolated bottom dissolved oxygen profile for September 2024.

### ChesMMAP November 2024 Dissolved Oxygen Profiles




Figure 94: Interpolated bottom dissolved oxygen profile for November 2024.

## Appendix II - History of ChesMMAP sampling design

Historically, the ChesMMAP sampling protocol included five 80-site surveys per year, one each in March, May, July, September and November. This general schedule was occasionally interrupted by funding shortfalls and/or logistical hurdles (e.g., vessel breakdowns). The *R/V Bay Eagle*, a 19.8 m aluminum hull, twin diesel vessel owned and operated by VIMS, served as the sampling platform for all cruises during this time period. Fishes (and select invertebrates) were collected using a 13.7 m (headrope length), two-bridle, four-seam bottom trawl manufactured by Reidar's Manufacturing Inc. of New Bedford, MA. The top belly, bottom belly, and side panels of the net are constructed of 15.2 cm stretch mesh (2.6 mm diameter twine), and the cod-end is constructed of 7.6 cm stretch mesh (1.6 mm diameter twine). The bridles (legs) of the net are 6.1 m and connected directly to 1.3 m x 0.8 m steel-V trawl doors weighing 71.8 kg each. The trawl net was deployed with a single-warp system using 9.5 mm diameter stainless steel main cable and a 37.6 m bridle constructed of 7.9 mm stainless steel wire rope.

The goal of each cruise was to sample 80 sites throughout the mainstem of Chesapeake Bay. Sampling sites were selected using a stratified random design. The Bay was stratified by dividing the mainstem into five regions of 30 latitudinal minutes each (the upper and lower regions being slightly smaller and larger than 30 minutes, respectively). Regions were numbered 1 through 5 from north to south. Regions 1-3 coincide with the Maryland portion of the Bay and regions 4-5 correspond with Virginia waters (note that due to the irregular state boundary it is possible that sites in the very southernmost portion of Region 3 may actually be in Virginia and likewise sites in the northernmost reaches of Region 4 may be north of the state border). Within each region, three depth strata ranging from 3.0 m-9.1 m, 9.1 m-15.2 m, and >15.2 m were defined. A grid of 1.9 km<sup>2</sup> cells was superimposed over the mainstem, where each cell represented a potential sampling location. The number of sites sampled in each region and in each stratum was proportional to the surface area of water represented. Sites were sampled without replacement and those north of Pooles Island (39° 17' N) have not been sampled since July 2002 due to repeated loss of trawl gear.

Tows were normally conducted in the same general direction as the tidal current, as pilot work conducted in November 2001 indicated that the survey gear performed most consistently when towed with the current rather than against the current. The net was generally deployed at a 4:1 scope, which refers to the cable length: water depth ratio. For shallow sites, however, the bridle wires were always fully deployed, implying that the scope ratio could be quite high in these situations. The target tow speed was 3.0 knots, but this occasionally varied depending on wind and tidal conditions. Based on data collected from the net monitoring gear, tow speed and scope were adjusted to ensure that the net maintained expected geometry. Tows were 20 minutes in duration, unless obstructions or other logistical issues forced a tow to be shortened; if the duration of a tow was at least 10 minutes, it was considered valid. Computer software was used to record data from the net monitoring gear (i.e., wingspread and headrope height) as well as a continuous GPS stream during each tow. On occasions when the monitoring gear failed or was not deployed, the trawl geometry was assumed to follow cruise averages and beginning and ending tow coordinates were recorded by hand from the vessel's GPS system.

In October 2018, VIMS took possession of its new research vessel, the *R/V Virginia*, a 28.3 m steel hull vessel with twin diesels tied to a single controllable-pitch propeller and a dynamic positioning system for station-holding. This vessel replaced the *R/V Bay Eagle* as the sampling platform for the ChesMMAP survey, and all future ChesMMAP sampling will occur on this new ship. In addition to the change in vessel, adjustments to the sampling gear were required. The new sampling gear is a

“200 x 12 cm” (200, 12 cm meshes at the front of the net, i.e. the “fishing circle”) bottom trawl rigged with a 3.8 cm cookie sweep and using Thyboron Type IV 44” trawl doors. The cod end is lined with a 2.5 cm knotless liner with an effective mesh size of approximately 1.6 cm. This sampling system is a one-half scale version of the net used by the North East Area Monitoring and Assessment Program (NEAMAP) and Northeast Fisheries Science Center (NEFSC) surveys and is approximately a twice-sized version of the net now being used by the VIMS Juvenile Fishes and Blue Crab Trawl Survey.

The sampling schedule changed in 2019 due to a combination of increased costs associated with the *R/V Virginia* and a decreasing budget, such that 3 sampling cruises could be conducted. However, in considering the annual pattern of fish abundances and in examining the subsets of the data used for the various species’ abundance indices, an alternative approach was implemented. In the early season (March) cruise none of the data from sampling in Virginia are used for any abundance indices. Likewise, in late season sampling (November), data for only one species in Maryland strata are used. Rather than settling for 3 full cruises we now sample in March, June, September and November, with the March and November trips sampling only in the upper (Maryland) and lower (Virginia) regions, respectively. While not ideal, we can still sample during the entire spring/summer/fall annual cycle.

Knowing that significant changes would be coming to the survey with the change in research vessel and sampling gear, survey stratification was evaluated. Analyses revealed that the prior design was over-stratified, with small numbers of samples coming from small strata but being over-represented in the design due to the criterion of sampling at least three sites from every stratum. Both the number of regions and the number of depth strata were reduced. The prior three regions corresponding to the Maryland portion of the Bay were condensed to two and similarly the number of depth strata in each region was reduced from three (3.0 m-9.1 m, 9.1 m-15.2 m, and > 15.2 m) to two ( $\leq 12.2$  m,  $> 12.2$  m). Thus, the total number of strata sampled during any cruise was reduced from 14 (there was no deep stratum in Region 1) to 8. Regions are now described as regions A (upper Maryland), B (lower Maryland), C (upper Virginia) and D (lower Virginia) and depth strata are similarly named A (shallow) and B (deep). While it may be somewhat confusing to use a similar labeling system for both the regions and depth strata, these conventions provide a clear distinction from the previous classifications.

## Appendix III - Additional species profiles

### Blue crab, *Callinectes sapidus*




Figure 95: Site-level estimates of biomass (kg 10,000 m<sup>-2</sup>) of male blue crabs in 2024.




Figure 96: Site-level estimates of biomass (kg 10,000 m<sup>-2</sup>) of female blue crabs in 2024.

Clearnose skate, *Rostroraja eglanteria*




Figure 97: Site-level estimates of biomass (kg 10,000 m<sup>-2</sup>) of clearnose skate in 2024.




Figure 98: Indices of abundance for clearnose skate by number and biomass, for all ages combined. The calibrated R/V Bay Eagle data are blue, while the R/V Virginia data are light yellow.